

Contents

How Do We Understand Life?	1
PART I: BIOLOGICAL MOLECULES	4
Chapter 1 From Genes to RNA and Proteins	5
Chapter 2 Nucleic Acid Structure	51
Chapter 3 Glycans and Lipids	91
Chapter 4 Protein Structure	131
Chapter 5 Evolutionary Variation in Proteins	191
PART II: ENERGY AND ENTROPY	238
Chapter 6 Energy and Intermolecular Forces	239
Chapter 7 Entropy	293
Chapter 8 Linking Energy and Entropy: The Boltzmann Distribution	341
PART III: FREE ENERGY	382
Chapter 9 Free Energy	383
Chapter 10 Chemical Potential and the Drive to Equilibrium	413
Chapter 11 Voltages and Free Energy	459
PART IV: MOLECULAR INTERACTIONS	530
Chapter 12 Molecular Recognition: The Thermodynamics of Binding	531
Chapter 13 Specificity of Macromolecular Recognition	581
Chapter 14 Allostery	633
PART V: KINETICS AND CATALYSIS	672
Chapter 15 The Rates of Molecular Processes	673
Chapter 16 Principles of Enzyme Catalysis	721
Chapter 17 Diffusion and Transport	787
PART VI: ASSEMBLY AND ACTIVITY	838
Chapter 18 Folding	839
Chapter 19 Fidelity in DNA and Protein Synthesis	887
Glossary	939
Index	965

Detailed Contents

How Do We Understand Life?	1	1.18 Splicing of RNA in eukaryotic cells can generate a diversity of RNAs from a single gene	39
PART I: BIOLOGICAL MOLECULES	4	1.19 The genetic code relates triplets of nucleotides in a gene sequence to each amino acid in a protein sequence	39
Chapter 1 From Genes to RNA and Proteins	5	1.20 Transfer RNAs work with the ribosome to translate mRNA sequences into proteins	42
A. INTERACTIONS BETWEEN MOLECULES	6	1.21 The mechanism for the transfer of genetic information is highly conserved	43
1.1 The energy of interaction between two molecules is determined by noncovalent interactions	6	1.22 The discovery of retroviruses showed that information stored in RNA can be transferred to DNA	44
1.2 Neutral atoms attract and repel each other at close distances through van der Waals interactions	8	Summary	46
1.3 Ionic interactions between charged atoms can be very strong, but are attenuated by water	10	Key Concepts	47
1.4 Hydrogen bonds are very common in biological macromolecules	12	Problems	48
B. INTRODUCTION TO NUCLEIC ACIDS AND PROTEINS	15	Further Reading	50
1.5 Nucleotides have pentose sugars attached to nitrogenous bases and phosphate groups	15	Chapter 2 Nucleic Acid Structure	51
1.6 The nucleotide bases in RNA and DNA are substituted pyrimidines or purines	18	A. DOUBLE-HELICAL STRUCTURES OF RNA AND DNA	52
1.7 DNA and RNA are formed by sequential reactions that utilize nucleotide triphosphates	20	2.1 The double helix is the principal secondary structure of DNA and RNA	52
1.8 DNA forms a double helix with antiparallel strands	22	2.2 Hydrogen bonding between bases is important for the formation of double helices, but its effect is weakened due to interactions with water	53
1.9 The double helix is stabilized by the stacking of base pairs	24	2.3 The electronic polarization of the bases contributes to strong stacking interactions between bases	54
1.10 Proteins are polymers of amino acids	25	2.4 Metal ions help shield electrostatic repulsions between the phosphate groups	55
1.11 Proteins are formed by connecting amino acids by peptide bonds	25	2.5 There are two common relative orientations of the base and the sugar	56
1.12 Amino acids are classified based on the properties of their sidechains	29	2.6 The ribose ring has alternate conformations defined by the sugar pucker	56
1.13 Proteins appear irregular in shape	30	2.7 RNA cannot adopt the standard Watson-Crick double-helical structure because of constraints on its sugar pucker	58
1.14 Protein chains fold up to form hydrophobic cores	31	2.8 The standard Watson-Crick model of double-helical DNA is the B-form	59
1.15 α helices and β sheets are the architectural elements of protein structure	31	2.9 B-form DNA allows sequence-specific recognition of the major groove, which has a greater information content than the minor groove	60
C. REPLICATION, TRANSCRIPTION, AND TRANSLATION	35	2.10 RNA adopts the A-form double-helical conformation	61
1.16 DNA replication is a complex process involving many protein machines	35	2.11 The major groove of A-form double helices is less accessible to proteins than that of B-form DNA	62
1.17 Transcription generates RNAs whose sequences are dictated by the sequence of nucleotides in genes	38		

2.12	Z-form DNA is a left-handed double-helical structure	62	B.	LIPIDS AND MEMBRANES	108
2.13	The DNA double helix is quite deformable	65	3.13	The most abundant lipids are glycerophospholipids	109
2.14	DNA supercoiling can occur when the ends of double helices are constrained	67	3.14	Other classes of lipids have different molecular frameworks	110
2.15	Writhe, linking number, and twist are mathematical parameters that describe the supercoiling of DNA	69	3.15	Lipids form organized structures spontaneously	113
2.16	The writhe, twist, and linking number are related to each other in a simple way	70	3.16	The shapes of lipid molecules affect the structures they form	113
2.17	The DNA in cells is supercoiled	71	3.17	Detergents are amphiphilic molecules that tend to form micelles rather than bilayers	115
2.18	Local conformational changes in the DNA also affect supercoiling	72	3.18	Lipids in bilayers move freely in two dimensions	116
B.	THE FUNCTIONAL VERSATILITY OF RNA	73	3.19	Lipid composition affects the physical properties of membranes	118
2.19	Wobble base pairs are often seen in RNA	73	3.20	Proteins can be associated with membranes by attachment to lipid anchors	121
2.20	Nonstandard base-pairing is common in RNA	75	3.21	Lipid molecules can be sequestered and transported by proteins	122
2.21	Some RNA molecules contain modified nucleotides	76	3.22	Different kinds of cells and organelles have different membrane compositions	123
2.22	A tetraloop is a common secondary structural motif that caps RNA hairpins	79	3.23	Cell walls are reinforced membranes	125
2.23	Interactions with metal ions help RNAs to fold	80	Summary		126
2.24	RNA tertiary structure involves interactions between secondary structural elements	81	Key Concepts		127
2.25	Helices in RNA often interact through coaxial base stacking or the formation of pseudoknots	82	Problems		128
2.26	Various interactions between nucleotides stabilize RNA tertiary structure	84	Further Reading		129
Summary		86			
Key Concepts		87			
Problems		88			
Further Reading		90			
Chapter 3 Glycans and Lipids	91		Chapter 4 Protein Structure	131	
A.	GLYCANS	91	A.	GENERAL PRINCIPLES	131
3.1	Simple sugars are comprised primarily of hydroxylated carbons	91	4.1	Protein structures display a hierarchical organization	131
3.2	Many cyclic sugar molecules can exist in alternative anomeric forms	92	4.2	Protein domains are the fundamental units of tertiary structure	133
3.3	Sugar rings often have many low energy conformations	94	4.3	Protein folding is driven by the formation of a hydrophobic core	134
3.4	Many sugars are structural isomers of identical composition, but with different stereochemistry	95	4.4	The formation of α helices and β sheets satisfies the hydrogen-bonding requirements of the protein backbone	136
3.5	Some sugars have other chemical functionalities in addition to alcohol groups	97	B.	BACKBONE CONFORMATION	137
3.6	Glycans form polymeric structures that can have branched linkages	98	4.5	Protein folding involves conformational changes in the peptide backbone	137
3.7	Differences in anomeric linkages lead to dramatic differences in polymeric forms of glucose	99	4.6	Amino acids are chiral and only the L form stereoisomer is found in genetically encoded proteins	138
3.8	Acetylation or other chemical modification leads to diversity in sugar polymer properties	101	4.7	The peptide bond has partial double bond character, so rotations about it are hindered	139
3.9	Glycans may be attached to proteins or lipids	102	4.8	Peptide groups can be in <i>cis</i> or <i>trans</i> conformations	140
3.10	The decoration of proteins with glycans is not templated	104	4.9	The backbone torsion angles ϕ (phi) and ψ (psi) determine the conformation of the protein chain	141
3.11	Glycan modifications alter the properties of proteins	105	4.10	The Ramachandran diagram defines the restrictions on backbone conformation	142
3.12	Protein-glycan interactions are important in cellular recognition	106	4.11	α helices and β strands are formed when consecutive residues adopt similar values of ϕ and ψ	143
			4.12	Loop segments have residues with very different values of ϕ and ψ	146
			4.13	α helices and β strands are often amphipathic	147
			4.14	Some amino acids are preferred over others in α helices	149

C.	STRUCTURAL MOTIFS AND DOMAINS IN SOLUBLE PROTEINS	150	4.42	Conformational changes in retinal impose directionality to proton flow in bacteriorhodopsin	181
4.15	Secondary structure elements are connected to form simple motifs	150	4.43	Active transporters cycle between conformations that are open to the interior or the exterior of the cell	183
4.16	Amphipathic α helices can form dimeric structures called coiled coils	153	4.44	ATP binding and hydrolysis provides the driving force for the transport of sugars into the cell	184
4.17	Hydrophobic sidechains in coiled coils are repeated in a heptad pattern	155		Summary	185
4.18	α helices that are integrated into complex protein structures do not usually form coiled coils	156		Key Concepts	187
4.19	The sidechains of α helices form ridges and grooves	157		Problems	188
4.20	α helices pack against each other with a limited set of crossing angles	157		Further Reading	189
4.21	Structures with alternating α helices and β strands are very common	159		Chapter 5 Evolutionary Variation in Proteins	191
4.22	α/β barrels occur in many different enzymes	161	A.	THE THERMODYNAMIC HYPOTHESIS	191
4.23	α/β open-sheet structures contain α helices on both sides of the β sheet	162	5.1	The structure of a protein is determined by its sequence	191
4.24	Proteins with antiparallel β sheets often form structures called β barrels	162	5.2	The thermodynamic hypothesis was first established for an enzyme known as ribonuclease-A, which can be unfolded and folded reversibly	192
4.25	Up-and-down β barrels have a simple topology	163	5.3	By counting the number of possible rearrangements of disulfide bonds, we can confirm that ribonuclease-A is completely unfolded by urea and reducing agents	194
4.26	Up-and-down β sheets can form repetitive structures	163	B.	SEQUENCE COMPARISONS AND THE BLOSUM MATRIX	195
4.27	Greek key motifs occur frequently in antiparallel β structures	164	5.4	Protein structure is conserved during evolution while amino acid sequences vary	195
4.28	Certain structural motifs can be repeated almost endlessly to form elongated structures	165	5.5	The globin fold is preserved in proteins that share very little sequence similarity	198
4.29	Catalytic sites are usually located within core elements of protein folds	167	5.6	Similarities in protein sequences can be quantified by considering the frequencies with which amino acids are substituted for each other in related proteins	201
4.30	Binding sites are often located at the interfaces between domains	168	5.7	The BLOSUM matrix is a commonly used set of amino acid substitution scores	201
D.	STRUCTURAL PRINCIPLES OF MEMBRANE PROTEINS	169	5.8	The first step in deriving substitution scores is to determine the frequencies of amino acid substitutions and correct for amino acid abundances	202
4.31	Lipid bilayers form barriers that are nearly impermeable to polar molecules	169	5.9	The substitution score is defined in terms of the logarithm of the substitution likelihood	204
4.32	Membrane proteins have distinct regions that interact with the lipid bilayer	170	5.10	The BLOSUM substitution scores reflect the chemical properties of the amino acids	207
4.33	The hydrophobicity of the lipid bilayer requires the formation of regular secondary structure within the membrane	171	5.11	Substitution scores are used to align sequences and to detect similarities between proteins	208
4.34	The more polar sidechains are rarely found within membrane-spanning α helices, except when they are required for specific functions	172	C.	STRUCTURAL VARIATION IN PROTEINS	209
4.35	Transmembrane α helices can be predicted from amino acid sequences	174	5.12	Small but significant differences in protein structures arise from differences in sequences	209
4.36	Hydrophobicity scales are used to identify transmembrane helices	175	5.13	Proteins retain a common structural core as their sequences diverge	210
4.37	Integral membrane proteins are stabilized by van der Waals contacts and hydrogen bonds	177	5.14	Structural overlap within the common core decreases as the sequences of proteins diverge	211
4.38	Porins contain β barrels that form transmembrane channels	178	5.15	Sequence comparisons alone are insufficient to establish structural similarity between distantly related proteins	212
4.39	Pumps and transporters use energy to move molecules across the membrane	179			
4.40	Bacteriorhodopsin uses light energy to pump protons across the membrane	180			
4.41	A hydrogen-bonded chain of water molecules can serve as a proton conducting "wire"	180			

5.16	The amino acids have preferences for different environments in folded proteins	213	6.7	The heat capacity of an ideal monatomic gas is constant	253
5.17	Fold-recognition algorithms evaluate the probability that the sequence of a protein is compatible with a known three-dimensional structure	214	6.8	The heat capacity of a macromolecular solution increases and then decreases with temperature	257
5.18	The 3D-1D profile method maps three-dimensional structural information onto a one-dimensional set of environmental descriptors	216	6.9	The potential energy of a molecular system is the energy stored in molecules and their interactions	259
5.19	The database of known protein structures is used to generate a scoring matrix that gives the likelihood of finding each amino acid in a particular environmental class	217	6.10	The Boltzmann distribution describes the population of molecules in different energy levels	261
5.20	The 3D-1D profile method matches sequences with structures	218	6.11	The energy required to break interatomic interactions in folded macromolecules gives rise to the peak in heat capacity	264
D.	THE EVOLUTION OF MODULAR DOMAINS	220	C.	ENERGETICS OF INTERMOLECULAR INTERACTIONS	265
5.21	Domains are the fundamental unit of protein evolution	220	6.12	Simplified energy functions are used to calculate molecular potential energies	265
5.22	Domains can be organized into families with similar folds	220	6.13	Empirical potential energy functions enable rapid calculation of molecular energies	266
5.23	The number of distinct fold families is likely to be limited	224	6.14	The energies of covalent bonds are approximated by functions such as the Morse potential	267
5.24	Protein domains are remarkably tolerant of changes in amino acid sequence, even in the hydrophobic core	225	6.15	Other terms in the energy function describe torsion angles and the deformations in the angles between covalent bonds	270
5.25	Structural plasticity in protein domains increases the tolerance to mutation	227	6.16	The van der Waals energy term describes weak attractions and strong repulsions between atoms	272
5.26	The Rossmann fold is found in many nucleotide binding proteins	228	6.17	Atoms in proteins and nucleic acids are partially charged	274
5.27	Thioredoxin reductase and glutathione reductase are enzymes that diverged from a common ancestor, but their active sites arose through convergent evolution	230	6.18	Electrostatic interactions are governed by Coulomb's law	275
Summary		232	6.19	Hydrogen bonds are an important class of electrostatic interactions	277
Key Concepts		234	6.20	Empirical energy functions are used in computer programs to calculate molecular energies	279
Problems		235	6.21	Interactions with water weaken the effective strengths of hydrogen bonds in proteins	281
Further Reading		237	6.22	The presence of hydrogen-bonding groups in a protein is important for solubility and specificity	282
PART II: ENERGY AND ENTROPY		238	6.23	The water surrounding protein molecules strongly influences electrostatic interactions	283
Chapter 6 Energy and Intermolecular Forces		239	6.24	The shapes of proteins change the electrostatic fields generated by charges within the protein	285
A.	THERMODYNAMICS OF HEAT TRANSFER	240	Summary		287
6.1	In order to keep track of changes in energy, we define the region of interest as the "system"	240	Key Concepts		288
6.2	Energy released by chemical reactions is converted to heat and work	242	Problems		289
6.3	The first law of thermodynamics states that energy is conserved	243	Further Reading		292
6.4	For a process occurring under constant pressure conditions, the heat transferred is equal to the change in the enthalpy of the system	246	Chapter 7 Entropy		293
6.5	Changes in energy do not always indicate the direction of spontaneous change	250	A.	COUNTING STATISTICS AND MULTIPLICITY	294
6.6	The isothermal expansion of an ideal gas occurs spontaneously even though the energy of the gas does not change	251	7.1	Different sequences of outcomes in a series of coin tosses have equal probabilities	294
B.	HEAT CAPACITIES AND THE BOLTZMANN DISTRIBUTION	253	7.2	When considering aggregate outcomes, the most likely result is the one that has maximum multiplicity	295
			7.3	The multiplicity of an outcome of coin tosses can be calculated using a simple formula involving factorials	297

7.4	The concept of multiplicity is broadly applicable in biology because a series of coin flips is analogous to a collection of molecules in alternative states	300
7.5	The binding of ligands to a receptor can be monitored by fluorescence microscopy	301
7.6	Pascal's triangle describes the multiplicity of outcomes for a series of binary events	302
7.7	The binomial distribution governs the probability of events with binary outcomes	304
7.8	When the number of events is large, Stirling's approximation simplifies the calculation of the multiplicity	306
7.9	The relative probability of two outcomes is given by the ratios of their multiplicities	307
7.10	As the number of events increases, the less likely outcomes become increasingly rare	308
7.11	For coin tosses, outcomes with equal numbers of heads and tails have maximal multiplicity	310
7.12	When the number of events is very large, the probability distribution is well approximated by a Gaussian distribution	311
7.13	The Gaussian distribution is centered at the mean value and has a width that is proportional to the standard deviation	312
7.14	Application of the Gaussian distribution enables statistical analysis of a series of binary outcomes	315
B.	ENTROPY	317
7.15	The logarithm of the multiplicity ($\ln W$) is related to the entropy	317
7.16	The multiplicity of a molecular system is the number of equivalent configurations of the molecules (microstates)	318
7.17	The multiplicity of a system increases as the volume increases	319
7.18	For a large number of atoms, the state with maximal multiplicity is the state that is observed at equilibrium	322
7.19	The Boltzmann constant, k_B , is a proportionality constant linking entropy to the logarithm of the multiplicity ($\ln W$)	325
7.20	The change in entropy is related to the heat transferred during a process	326
7.21	The work done in a near-equilibrium process is greater than for a nonequilibrium process	327
7.22	The work done in a near-equilibrium process is related to the change in entropy	329
7.23	The statistical and thermodynamic definitions of entropy are equivalent	330
7.24	The second law of thermodynamics states that spontaneous change occurs in the direction of increasing entropy	331
7.25	Diffusion across a semipermeable membrane can lead to unequal numbers of molecules on the two sides of the membrane	332
Summary		335
Key Concepts		336
Problems		337
Further Reading		339

Chapter 8 Linking Energy and Entropy: The Boltzmann Distribution 341

A.	ENERGY DISTRIBUTIONS AND ENTROPY	341
8.1	The thermodynamic definition of the entropy provides a link to experimental observations	341
8.2	The concept of temperature provides a connection between the statistical and thermodynamic definitions of entropy	343
8.3	Energy distributions describe the populations of molecules with different energies	344
8.4	The multiplicity of an energy distribution is the number of equivalent configurations of molecules that results in the same energy distribution	344
8.5	The multiplicity of a system with different energy levels can be calculated by counting the number of equivalent molecular rearrangements of energy	347
B.	THE BOLTZMANN DISTRIBUTION	350
8.6	For large numbers of molecules, a probabilistic expression for the entropy is more convenient	350
8.7	The multiplicity of a system changes when energy is transferred between systems	354
8.8	Systems in thermal contact exchange heat until the combined entropy of the two systems is maximal	356
8.9	Many energy distributions are consistent with the total energy of a system, but some have higher multiplicity than others	359
8.10	The energy distribution at equilibrium must have an exponential form	360
8.11	The partition function indicates the accessibility of the higher energy levels of the system	363
8.12	For large numbers of molecules, non-Boltzmann distributions of the energy are highly unlikely	367
C.	ENTROPY AND TEMPERATURE	368
8.13	The rate of change of entropy with respect to energy is related to the temperature	368
8.14	The statistical and thermodynamic definitions of the entropy are equivalent	375
Summary		377
Key Concepts		378
Problems		379
Further Reading		381

PART III: FREE ENERGY 382

Chapter 9 Free Energy 383

A.	FREE ENERGY	384
9.1	The combined entropy of the system and the surroundings increases for a spontaneous process	384
9.2	The change in entropy of the surroundings is related to the change in energy and volume of the system	386
9.3	The Gibbs free energy (G) of the system always decreases in a spontaneous process occurring at constant pressure and temperature	387

9.4	The Helmholtz free energy (A) determines the direction of spontaneous change when the volume is constant	389	10.6	The chemical potentials of the reactants and products are balanced at equilibrium	422
B.	STANDARD FREE-ENERGY CHANGES	390	10.7	The concentrations of reactants and products at equilibrium define the equilibrium constant (K), which is related to the standard free energy change (ΔG°) for the reaction	424
9.5	Standard free-energy changes are defined with reference to defined standard states	390	10.8	Equilibrium constants can be used to calculate the extent of reaction at equilibrium	425
9.6	The zero point of the free-energy scale is set by the free energy of the elements in their most stable forms	391	10.9	The free-energy change for the reaction (ΔG), not the standard free-energy change (ΔG°), determines the direction of spontaneous change	426
9.7	Thermodynamic cycles allow the determination of the free energies of formation of complex molecules from simpler ones	392	10.10	The ratio of the reaction quotient (Q) to the equilibrium constant (K) determines the thermodynamic drive of a reaction	427
9.8	The free energy of formation of glucose is obtained by considering three combustion reactions	394	10.11	ATP concentrations are maintained at high levels in cells, thereby increasing the driving force for ATP hydrolysis	427
9.9	Enthalpies and entropies of formation can be combined to give the free energy of formation	395	C.	ACID-BASE EQUILIBRIA	428
9.10	Calorimetric measurements yield the standard enthalpy changes associated with combustion reactions	396	10.12	The Henderson-Hasselbalch equation relates the pH of a solution of a weak acid to the concentrations of the acid and its conjugate base	429
9.11	The entropy of formation of a compound is derived from heat capacity measurements	396	10.13	The proton concentration ($[H^+]$) in pure water at room temperature corresponds to a pH value of 7.0	430
C.	FREE ENERGY AND WORK	398	10.14	The temperature dependence of the equilibrium constant allows us to determine the values of ΔH° and ΔS°	431
9.12	Expansion work is not the only kind of work that can be done by a system	398	10.15	Weak acids, such as acetic acid, dissociate very little in water	432
9.13	Chemical work involves changes in the numbers of molecules	400	10.16	Solutions of weak acids and their conjugate bases act as buffers	433
9.14	The decrease in the Gibbs free energy for a process is the maximum amount of non-expansion work that the system is capable of doing under constant pressure and temperature	400	10.17	The charges on biological macromolecules are affected by the pH	435
9.15	The coupling of ATP hydrolysis to work underlies many processes in biology	402	10.18	The charge on an amino acid sidechain can be altered by interactions in the folded protein	436
9.16	The synthesis of ATP is coupled to the movement of ions across the membrane, down a concentration gradient	405	D.	FREE-ENERGY CHANGES IN PROTEIN FOLDING	438
Summary		408	10.19	The protein folding reaction is simplified by ignoring intermediate conformations	438
Key Concepts		409	10.20	Protein folding results from a balance between energy and entropy	439
Problems		409	10.21	The entropy of the unfolded protein chain is proportional to the logarithm of the number of conformations of the chain	440
Further Reading		411	10.22	The number of conformations of the unfolded chain can be estimated by counting the number of low-energy torsional isomers	442
Chapter 10 Chemical Potential and the Drive to Equilibrium	413		10.23	The free-energy change opposes protein folding if the entropy of water molecules is not considered	443
A.	CHEMICAL POTENTIAL	413	10.24	Protein folding is driven by an increase in water entropy	444
10.1	The chemical potential of a molecular species is the molar free energy of that species	414	10.25	Calorimetric measurements allow the experimental determination of the free energy of protein folding	446
10.2	Molecules move spontaneously from regions of high chemical potential to regions of low chemical potential	414	10.26	The heat capacity of a protein solution depends on the relative population of folded and unfolded molecules, and on the energy required to unfold the protein	446
10.3	Biochemical reactions are assumed to occur in ideal and dilute solutions, which simplifies the calculation of the chemical potential	416			
10.4	The chemical potential is proportional to the logarithm of the concentration	417			
10.5	Chemical potentials at arbitrary concentrations are calculated with reference to standard concentrations	421			
B.	EQUILIBRIUM CONSTANTS	422			

10.27	The area under the peak in the melting curve is the enthalpy change for unfolding at the melting temperature	448	11.17	An electrical potential difference across the membrane is essential for the functioning of all cells	484
10.28	The heat capacities of the folded and unfolded protein allow the determination of ΔH° and ΔS° for unfolding at any temperature	449	11.18	The sodium-potassium pump hydrolyzes ATP to move Na^+ ions out of the cell with the coupled movement of K^+ ions into the cell	486
10.29	Folded proteins become unstable at very low temperature because of changes in ΔH° and ΔS°	452	11.19	Sodium and potassium channels allow ions to move quickly across the membrane	487
Summary		453	11.20	Sodium and potassium channels contain a conserved tetrameric pore domain	489
Key Concepts		455	11.21	A large vestibule within the channel reduces the distance over which ions have to move without associated water molecules	490
Problems		456	11.22	Carbonyl groups in the selectivity filter provide specificity for K^+ ions by substituting for the inner-sphere waters	491
Further Reading		457	11.23	Rapid transit of K^+ ions through the channel is facilitated by hopping between isoenergetic binding sites	492
Chapter 11 Voltages and Free Energy	459		D. THE TRANSMISSION OF ACTION POTENTIALS IN NEURONS	493	
A. OXIDATION-REDUCTION REACTIONS IN BIOLOGY	459		11.24	The asymmetric distribution of ions across the cell membrane generates an equilibrium membrane potential	493
11.1	Reactions involving the transfer of electrons are referred to as oxidation-reduction reactions	459	11.25	The Nernst equation relates the equilibrium membrane potential to the concentrations of ions inside and outside the cell	494
11.2	Biologically important redox-active metals are bound to proteins	460	11.26	Cell membranes act as electrical capacitors	496
11.3	Nicotinamide adenine dinucleotide (NAD^+) is an important mediator of redox reactions in biology	460	11.27	The depolarization of the membrane is a key step in initiating a neuronal signal	498
11.4	Flavins and quinones can undergo oxidation or reduction in two steps of one electron each	461	11.28	Membrane potentials are altered by the movement of relatively few ions, enabling rapid axonal transmission	499
11.5	The oxidation of glucose is coupled to the generation of NADH and FADH_2	463	11.29	The propagation of voltage changes can be understood by treating the axon as an electrical circuit	500
11.6	Mitochondria are cellular compartments in which NADH and FADH_2 are used to generate ATP	465	11.30	The propagation of changes in membrane potential in the axon are described by the cable equation	501
11.7	Absorption of light creates molecules with high reducing power in photosynthesis	467	11.31	The resting membrane potential is determined by a combination of the basal conductances of potassium and sodium channels	505
B. REDUCTION POTENTIALS AND FREE ENERGY	469		11.32	The propagation of a voltage spike without triggering voltage-gated ion channels is known as passive spread	506
11.8	Electrochemical cells can be constructed by linking two redox couples	470	11.33	If membrane currents are neglected, then the cable equation is analogous to a diffusion equation	507
11.9	The voltage generated by an electrochemical cell with the reactants at standard conditions is known as the standard cell potential	473	11.34	Leakage through open ion channels limits the spread of a voltage perturbation	509
11.10	The electric potential difference (voltage) between two points is the work done in moving a unit charge between the two points	474	11.35	The time taken to develop a membrane potential is determined by the conductance of the membrane and its capacitance	510
11.11	Standard reduction potentials are related to the standard free-energy change of the redox reaction underlying the electrochemical cell	475	11.36	Myelination of mammalian neurons facilitates the transmission of action potentials	513
11.12	Electrode potentials are measured relative to a standard hydrogen electrode	477	11.37	Action potentials are regenerated periodically as they travel down the axon	514
11.13	Tabulated values of standard electrode potentials allow ready calculation of the standard potential of an electrochemical cell	478	11.38	A positively charged sensor in voltage-gated ion channels moves across the membrane upon depolarization	517
11.14	The Nernst equation describes how the potential changes with the concentrations of the redox reactants	480			
11.15	The standard state for reduction potentials in biochemistry is pH 7	480			
C. ION PUMPS AND CHANNELS IN NEURONS	481				
11.16	Neuronal cells use electrical signals to transmit information	482			

11.39	The structures of voltage-gated K^+ channels show that the voltage sensors form paddle-like structures that surround the core of the channel	520	12.16	Induced-fit binding occurs through selection by the ligand of one among many preexisting conformations of the protein	557
11.40	The crystal structure of a voltage-gated K^+ channel suggests how the voltage sensor opens and closes the channel	521	12.17	Conformational changes in the protein underlie the specificity of a cancer drug known as imatinib	559
Summary		524	12.18	Conformational changes in the target protein can weaken the affinity of an inhibitor	560
Key Concepts		525	12.19	The strength of noncovalent interactions usually correlates with hydrophobic interactions	562
Problems		526	12.20	Cholesterol-lowering drugs known as statins take advantage of hydrophobic interactions to block their target enzyme	563
Further Reading		527	12.21	The apparent affinity of a competitive inhibitor for a protein is reduced by the presence of the natural ligand	566
PART IV: MOLECULAR INTERACTIONS	530		12.22	Entropy lost by drug molecules upon binding is regained through the hydrophobic effect and the release of protein-bound water molecules	569
Chapter 12 Molecular Recognition: The Thermodynamics of Binding	531		12.23	Isothermal titration calorimetry allows us to determine the enthalpic and entropic components of the binding free energy	573
A. THERMODYNAMICS OF MOLECULAR INTERACTIONS	531		Summary		576
12.1	The affinity of a protein for a ligand is characterized by the dissociation constant, K_D	533	Key Concepts		578
12.2	The value of K_D corresponds to the concentration of free ligand at which the protein is half saturated	535	Problems		578
12.3	The dissociation constant is a dimensionless number, but is commonly referred to in concentration units	537	Further Reading		580
12.4	Dissociation constants are determined experimentally using binding assays	537	Chapter 13 Specificity of Macromolecular Recognition	581	
12.5	Binding isotherms plotted with logarithmic axes are commonly used to determine the dissociation constant	540	A. AFFINITY AND SPECIFICITY	581	
12.6	When the ligand is in great excess over the protein, the free ligand concentration, $[L]$, is essentially equal to the total ligand concentration	542	13.1	Both affinity and specificity are important in intermolecular interactions	581
12.7	Scatchard analysis makes it possible to estimate the value of K_D when the concentration of the receptor is unknown	543	13.2	Proteins often have to choose between several closely related targets	582
12.8	Scatchard analysis can be applied to unpurified proteins	544	13.3	Specificity is defined in terms of ratios of dissociation constants	584
12.9	Saturable binding is a hallmark of specific binding interactions	546	13.4	The specificity of binding depends on the concentration of ligand	585
12.10	The value of the dissociation constant, K_D , defines the ligand concentration range over which the protein switches from unbound to bound	546	13.5	Fractional occupancy and specificity are important for activities resulting from binding	587
12.11	The dissociation constant for a physiological ligand is usually close to the natural concentration of the ligand	548	13.6	Most macromolecular interactions are a compromise between affinity and specificity	587
B. DRUG BINDING BY PROTEINS	549		13.7	Fibroblast growth factors vary considerably in their affinities for receptors	588
12.12	Most drugs are developed by optimizing the inhibition of protein targets	549	13.8	The recognition of DNA by transcription factors involves discrimination between a very large numbers of off-target binding sites	590
12.13	Signaling molecules are protein targets in cancer drug development	549	13.9	Lowering the affinity of lac repressor for the operator switches on transcription	591
12.14	Most small molecule drugs work by displacing a natural ligand for a protein	552	B. PROTEIN-PROTEIN INTERACTIONS	593	
12.15	The binding of drugs to their target proteins often results in conformational changes in the protein	556	13.10	Protein-protein complexes involve interfaces between two folded domains or between a domain and a peptide segment	593
			13.11	SH2 domains are specific for peptides containing phosphotyrosine	595
			13.12	Individual SH2 domains cannot discriminate sharply between different phosphotyrosine-containing sequences	596
			13.13	Combinations of peptide recognition domains have higher specificity than individual domains	597

16.12 Substrate-dependent noncompetitive inhibitors only bind to the enzyme when the substrate is present	741	Further Reading	785
16.13 Some noncompetitive inhibitors are linked irreversibly to the enzyme	742	Chapter 17 Diffusion and Transport	787
16.14 In a ping-pong mechanism the enzyme becomes modified temporarily during the reaction	744	A. RANDOM WALKS	787
16.15 For a reaction with multiple substrates, the order of binding can be random or sequential	744	17.1 Microscopic motion is well described by trajectories called random walks	787
16.16 Enzymes with multiple binding sites can display allosteric (cooperative) behavior	746	17.2 The analysis of bacterial movement is simplified by considering one-dimensional random walks with uniform step lengths and time intervals	788
16.17 Product inhibition is a mechanism for regulating metabolite levels in cells	749	17.3 The probability distribution for the number of moves in one direction is given by a Gaussian function	789
C. PROTEIN ENZYMES	749	17.4 The probability of moving a certain distance in a one-dimensional random walk is also given by a Gaussian function	791
16.18 Enzymes can accelerate reactions by large amounts	750	17.5 The width of the distribution of displacements increases with the square root of time for random walks	794
16.19 Transition state stabilization is a major contributor to rate enhancement by enzymes	751	17.6 Random walks in two dimensions can be analyzed by combining two orthogonal one-dimensional random walks	796
16.20 Enzymes can act as acids or bases to enhance reaction rates	754	17.7 A two-dimensional random walk is described by two one-dimensional walks, but the effective step size for each is smaller by a factor of $\sqrt{2}$	798
16.21 Proximity effects are important for many reactions	756	17.8 The assumption of uniform step lengths along each axis means that the random walk occurs on a grid	798
16.22 The serine proteases are a large family of enzymes that contain a conserved Ser-His-Asp catalytic triad	758	17.9 A three-dimensional random walk is described by three orthogonal one-dimensional walks, and the effective step size for each is smaller by a factor of $\sqrt{3}$	801
16.23 Sidechain recognition positions the catalytic triad next to the peptide bond that is cleaved	758	17.10 The movement of bacteria in the presence of attractants or repellents is described by biased random walks	801
16.24 The specificities of serine proteases vary considerably, but the catalytic triad is conserved	760	B. MACROSCOPIC DESCRIPTION OF DIFFUSION	802
16.25 Peptide cleavage in serine proteases proceeds via a ping-pong mechanism	761	17.11 Fick's first law states that the flux of molecules is proportional to the concentration gradient	802
16.26 Angiotensin-converting enzyme is a zinc-containing protease that is an important drug target	763	17.12 Fick's second law describes the rate of change in concentration with time	804
16.27 Creatine kinase catalyzes phosphate transfer by stabilizing a planar phosphate intermediate	764	17.13 Integration of the diffusion equation allows us to calculate the change in concentration with time	805
16.28 Some enzymes work by populating disfavored conformations	766	17.14 The diffusion constant is related to the mean square displacement of molecules	807
16.29 Antibodies that bind transition state analogs can have catalytic activity	768	17.15 Diffusion constants depend on molecular properties such as size and shape	809
D. RNA ENZYMES	769	17.16 The diffusion constant is inversely related to the friction factor	810
16.30 Small self-cleaving ribozymes and ribonuclease proteins catalyze the same reaction	769	17.17 Viscosity is a measure of the resistance to flow	811
16.31 Self-cleaving ribozymes use nucleotide bases for catalysis, even though these do not have pK_a values well suited for proton transfer	769	17.18 Liquids with strong interactions between molecules have high viscosity	812
16.32 Hairpin ribozymes optimize hydrogen bonds to the transition state rather than to the initial or final states	771	17.19 The Stokes-Einstein equation allows us to calculate the diffusion coefficients of molecules	812
16.33 There are at least two possible mechanisms for bond cleavage by the hairpin ribozyme	773	17.20 The diffusion constants for nonspherical molecules are only slightly different from those calculated from the spherical approximation	814
16.34 The splicing reaction catalyzed by group I introns occurs in two steps	774	17.21 Diffusion-limited reaction rate constants can be calculated from the diffusion constants of molecules	815
16.35 Metal ions facilitate catalysis by group I introns	777		
16.36 Substitution of oxygen by sulfur in RNA helps identify metals that participate in catalysis	777		
Summary	780		
Key Concepts	781		
Problems	782		

17.22	One-dimensional searches on DNA increase the rate at which transcription factors find their targets	817	18.10	Changes in the sequence of a protein at certain positions can affect folding rates substantially	850
17.23	Restricting diffusion to two-dimensional membranes can slow down the rate of encounter but still speed up reactions	819	18.11	The nature of the transition state can be identified by mapping the effect of mutations on the folding and unfolding rates	852
17.24	Concentration gradients determine the outcomes of many biological processes	822	18.12	The process of protein folding can be described as funneled movement on a multidimensional free-energy landscape	856
17.25	Cells use motor proteins to transport cargo over long distances and to specific locations	823	B.	CHAPERONES FOR PROTEIN FOLDING	857
17.26	Vesicles are transported by kinesin motors that move along microtubule tracks	823	18.13	Many proteins tend to aggregate rather than fold	857
17.27	ATP hydrolysis provides a powerful driving force for kinesin movement	825	18.14	The high concentration of macromolecules inside the cell makes the problem of aggregation particularly acute	858
C.	EXPERIMENTAL MEASUREMENT OF DIFFUSION	826	18.15	Proteins inside the cell usually fold into a functional form rapidly	860
17.28	Diffusion constants can be measured experimentally in several ways	826	18.16	Some proteins form irreversible aggregates that are toxic to cells	861
17.29	Movement of molecules in solution can be driven by centrifugal forces	827	18.17	Molecular chaperones are proteins that prevent protein aggregation	863
17.30	Equilibrium centrifugation can be used to determine molecular weights	829	18.18	Hsp70 recognizes short peptides with sequences that are characteristic of the interior segments of proteins	866
17.31	Electrophoresis provides an alternative method for driving molecular motion	830	18.19	Hsp70 binds and releases protein chains in a cycle that is coupled to ATP binding and hydrolysis	866
17.32	The electrophoretic mobility of nucleic acids decreases with size	831	18.20	The GroEL chaperonin forms a hollow double-ring structure within which protein molecules can fold	868
17.33	Gel electrophoresis analysis of proteins is useful for size determination	832	18.21	GroEL works like a two-stroke engine, binding and releasing proteins	870
Summary		833	18.22	GroEL-GroES can accelerate the folding of proteins through passive and active mechanisms	872
Key Concepts		834	C.	RNA FOLDING	872
Problems		835	18.23	The electrostatic field around RNA leads to the diffuse localization of metal ions	873
Further Reading		836	18.24	RNA folding can be driven by increasing the concentration of metal ions	874
PART VI: ASSEMBLY AND ACTIVITY		838	18.25	RNAs form stable secondary structural elements, which increases their tendency to misfold	875
Chapter 18 Folding		839	18.26	RNA folding is hierarchical with multiple stable intermediates	877
A.	HOW PROTEINS FOLD	840	18.27	Collapse is an early event in the folding of RNA	878
18.1	Protein folding is governed by thermodynamics	840	18.28	RNA folding landscapes are highly rugged	880
18.2	The reversibility of protein folding can also be demonstrated by manipulating single molecules	841	Summary		882
18.3	Unfolded states of proteins correspond to wide distributions of different conformations	842	Key Concepts		883
18.4	Protein folding cannot be explained by an exhaustive search of conformational space	844	Problems		884
18.5	Many small proteins populate only fully unfolded and fully folded states	844	Further Reading		886
18.6	The order in which secondary and tertiary interactions form can vary in different proteins	845	Chapter 19 Fidelity in DNA and Protein Synthesis		887
18.7	Folding rates are faster when residues close in sequence end up close together in the folded structure	846	A.	MEASURING THE STABILITY OF DNA DUPLEXES	888
18.8	The folding of some proteins involves the formation of transiently stable intermediates	847	19.1	The difference in free energy between matched and mismatched base pairs can be determined by measuring the melting temperature of DNA	888
18.9	Folding pathways can have multiple intermediates	850			

19.2	DNA melting can be studied by UV absorption spectroscopy	889	19.19	DNA polymerases recognize DNA using the backbone and minor groove	915
19.3	The changes in enthalpy and entropy associated with DNA melting can be determined from the concentration dependence of melting curves	890	19.20	DNA polymerases sense the shapes of correctly paired bases	917
19.4	DNA duplexes containing a mismatched base pair at one end are only marginally less stable than duplexes with matched bases	892	19.21	The shape of a nucleotide is more important for its being incorporated into DNA than its ability to form hydrogen bonds	918
19.5	The entropy of each DNA chain is reduced upon forming a duplex	894	19.22	The growing DNA strand can shuttle between the polymerase and exonuclease active sites	919
19.6	The stability of DNA depends on the pattern on base stacks in the duplex	895	C.	HOW RIBOSOMES ACHIEVE FIDELITY	920
19.7	Base stacking is more important than hydrogen bonding in determining the stability of DNA helices	897	19.23	The ribosome has two subunits, each of which is a large complex of RNA and proteins	921
B.	FIDELITY IN DNA REPLICATION	898	19.24	Protein synthesis on the ribosome occurs as a repeated series of steps of tRNA and protein binding, with conformational changes in the ribosome	921
19.8	The process of DNA replication is very accurate	898	19.25	Selection of the correct A-site tRNA by base-pairing alone cannot explain ribosome fidelity	923
19.9	The energy of DNA base-pairing cannot explain the accuracy of DNA replication	900	19.26	A ribosome-induced bend in the EF-Tu•tRNA complex plays an important role in generating specificity	924
19.10	The overall process of DNA synthesis can be described as a series of kinetic steps	902	19.27	The ribosome undergoes conformational changes during the process of tRNA selection	925
19.11	Primer elongation by polymerase is quite rapid	904	19.28	Tight interactions in the decoding center can only occur for correct codon-anticodon pairs	926
19.12	The rate-limiting step in the DNA synthesis reaction is a conformational change in DNA polymerase	905	19.29	Coupling of the decoding center and the GTPase active site of EF-Tu involves multiple conformational rearrangements	929
19.13	Determination of the values of $V_{\max}$ and K_M for the incorporation of correct and incorrect base pairs provides insights into fidelity	907	19.30	The active site of EF-Tu needs only a small rearrangement to be activated	930
19.14	DNA polymerase has a nuclease activity that can remove bases from the 3' end of a DNA strand	908	19.31	Release of EF-Tu allows the aminoacyl group of the A-site tRNA to move to the peptidyl transfer center	931
19.15	The structure of DNA polymerase has fingers, palm, and thumb subdomains	909	19.32	The ribosome catalyzes peptidyl transfer	932
19.16	DNA polymerase binds DNA using the "palm" and nearly encircles it	910	Summary		934
19.17	The active site of polymerase contains two metals ions that catalyze nucleotide addition	911	Key Concepts		935
19.18	A conformational change in DNA polymerase upon binding dNTP contributes to replication fidelity	913	Problems		936
			Further Reading		937