

Contents in Detail

Preface	vii	Transcriptional control can operate at the level of chromatin structure and at the level of production of the primary RNA transcript	20
Acknowledgments	viii		
Online resources	ix		
1 Levels of Gene Control	1	1.5 Regulatory RNAs and the regulation of gene expression	21
Introduction	1	RNA molecules of different lengths are involved in the control of gene expression	21
1.1 The protein content of different cell types is different	1	MicroRNAs are processed from a single-stranded precursor which folds to form a double-stranded hairpin loop	21
Specific methods can be used to study the expression of individual proteins in tissues and cells	1	Many small interfering RNAs are processed from a double-stranded precursor	24
General methods can be used for studying the overall protein composition of tissues and cells	2	Piwi-interacting RNAs are processed in a Dicer-independent manner	26
1.2 The mRNA content of different cell types is different	5	Small RNAs can move from cell to cell and from one part of an organism to another	28
Specific methods can be used to study the expression of individual mRNAs in different tissues and cells	5	Long noncoding RNAs function by a variety of mechanisms	28
General methods can be used to study the overall population of mRNAs expressed in different tissues and cells	7	Conclusions	32
1.3 The DNA content of different cell types is generally the same	9	Key concepts	33
Specific methods can be used to study individual genes in different tissues and cells	9	Further reading	33
General methods can be used to study the total DNA in different tissues and cells	10	2 Structure of Chromatin	35
Exceptional cases do exist in which changes to the DNA occur in specific tissues or cell types	13	Introduction	35
1.4 Transcriptional or post-transcriptional control?	15	Regulation of transcription in eukaryotes is much more complex than in prokaryotes	35
Pulse labeling studies directly demonstrate transcriptional control	16	2.1 Commitment to the differentiated state and its stability	36
Nuclear run-on assays allow transcriptional control to be demonstrated for a wide range of genes	17	Cells can remain committed to a particular differentiated state even in the absence of its phenotypic characteristics	36
Polytene chromosomes provide further evidence for transcriptional control	19	Cells can become committed to a particular differentiated state before actual phenotypic differentiation	37
		2.2 The nucleosome	39
		The nucleosome is the basic unit of chromatin structure	39

Nucleosome structure or position can be altered by chromatin remodeling processes	42	3.2 Alterations in DNA methylation in active or potentially active genes	71
2.3 Histone modifications and histone variants	46	Decreased DNA methylation is associated with active or potentially active genes	71
Histones are subject to a variety of post-translational modifications	46	DNA methylation plays a key role in regulating chromatin structure	73
Histone variants are encoded by distinct genes to those encoding the standard histone isoforms	52	DNA methylation patterns can be propagated stably through cell divisions	75
2.4 The 30 NM chromatin fiber	54	DNA methylation recruits inhibitory proteins that produce a tightly packed chromatin structure	78
The 30 nm fiber represents a further compaction of the beads-on-a-string structure	54	3.3 Modification of histones in the chromatin of active or potentially active genes	79
Histone H1 and post-translational modifications of the other histones are involved in the formation of the 30 nm fiber	55	3.4 Functional interaction of different histone modifications, DNA methylation, and regulatory RNAs	89
2.5 Structural and functional domains in chromatin	57	The different histone modifications functionally interact with one another	89
The 30 nm fiber is further compacted by looping	57	Histone modifications interact with DNA methylation to regulate chromatin structure	90
Locus control regions regulate the chromatin structure of a large region of DNA	59	Regulatory RNAs can induce alterations in chromatin structure	91
Insulators block the inappropriate spread of particular chromatin structures	61	3.5 Changes in chromatin structure in the regulatory regions of active or potentially active genes	95
Heterochromatin is a very tightly packed form of chromatin	62	DNaseI hypersensitive sites can be identified in active or potentially active genes	95
The chromosome is the visible result of chromatin compaction	62	DNaseI hypersensitive sites frequently correspond to regulatory DNA sequences	97
Conclusions	64	DNaseI hypersensitive sites represent areas which are either nucleosome-free or have an altered nucleosomal structure	98
Key concepts	65	Chromatin remodeling can be produced by proteins capable of displacing nucleosomes or altering their structure	99
Further reading	65	The SWI/SNF and NURF chromatin remodeling complexes are recruited to the DNA by a variety of different mechanisms	99
3 The Epigenome: Role of Chromatin Structure in Gene Control	67	The initial opening of tightly packed chromatin can involve the binding of pioneer transcription factors	102
Introduction	67		
3.1 Changes in chromatin structure in active or potentially active genes	68		
Active DNA is organized in a nucleosomal structure	68		
Active or potentially active chromatin shows enhanced sensitivity to DNaseI digestion	69		

3.6 Other situations in which chromatin structure is regulated	103		
In female mammals one of the two X chromosomes is inactivated	103		
The active and inactive X chromosomes have a different chromatin structure	104		
The XIST regulatory RNA is specifically transcribed on the inactive X chromosome	104		
Genomic imprinting involves the specific inactivation of either the maternally or paternally inherited copy of specific genes	106		
Imprinting involves changes in chromatin structure	107		
Conclusions	110		
Key concepts	111		
Further reading	112		
4 The Process of Transcription	115		
Introduction	115		
4.1 Transcription by RNA polymerases	115		
Transcription by RNA polymerase I is relatively simple	116		
Transcription by RNA polymerase III is more complex than for RNA polymerase I	117		
Transcription by RNA polymerase II is much more complex than transcription by RNA polymerases I and III	119		
Plants contain two additional RNA polymerases which are related to RNA polymerase II and play a key role in gene silencing by small RNAs	122		
Transcription by RNA polymerases I, II, and III has a number of common features	123		
Transcription takes place in defined regions of the nucleus	127		
4.2 Transcriptional elongation and termination	129		
Transcriptional elongation requires further phosphorylation of RNA polymerase II	129		
		Polymerase pausing is widespread and represents a potential control point for the regulation of transcription	130
		Termination of transcription occurs downstream of the polyadenylation signal	131
		4.3 The gene promoter	134
		The 70 kDa heat-shock protein gene contains a typical promoter for RNA polymerase II	135
		The <i>hsp70</i> gene promoter contains several DNA sequence motifs which are found in a variety of other gene promoters	135
		The heat-shock element is found only in heat-inducible genes	136
		Other response elements are found in the promoters of genes with different patterns of expression	138
		The proteins binding to short DNA sequence elements can be characterized by a variety of techniques	140
		The use of high-throughput methods such as ChIP-chip and ChIP-seq allows the analysis of DNA-binding sites and epigenetic modifications on a whole genome basis	144
		Promoter regulatory elements act by binding factors which either affect chromatin structure and/or influence transcription directly	145
		4.4 Enhancers and silencers	146
		Enhancers are regulatory sequences that act at a distance to increase gene expression	146
		Many enhancers have cell type- or tissue-specific activity	148
		Enhancers influence transcription both by recruiting DNA-binding transcription factors and by altering chromatin structure	150
		The "at a distance" action of enhancers involves both DNA-binding transcription factors and alteration of chromatin structure	152
		Many enhancers are transcribed to produce long noncoding RNAs	153

Silencers can act at a distance to inhibit gene expression	154	Several different classes of activation domain exist	183
Conclusions	155	Activators can interact with the TAFs within the TFIID complex	184
Key concepts	156	Activators can interact with co-activators	186
Further reading	157	Activators can interact with modulators of chromatin structure	187
5 Transcription Factors and Transcriptional Control	159	Activators have a multitude of targets	188
Introduction	159	5.3 Repression of transcription	190
5.1 DNA binding by transcription factors	162	Repressors can act indirectly by inhibiting the positive effect of activators	191
The helix-turn-helix motif is found in a number of transcription factors which regulate gene expression during embryonic development	162	Repressors can act directly by inhibiting the assembly or activity of the basal transcriptional complex	191
The helix-turn-helix domain found in homeodomain proteins is a DNA-binding domain	164	5.4 Regulation at transcriptional elongation	194
In the POU domain transcription factors, the homeodomain forms part of a larger DNA-binding motif	166	Regulation of transcription can occur at the elongation stage, as well as at initiation	194
The two-cysteine-two-histidine zinc finger is found in multiple copies in many transcription factors	168	Factors which regulate transcriptional elongation target the C-terminal domain of the RPB1 subunit of RNA polymerase II	197
The nuclear receptors contain two copies of a multi-cysteine zinc finger distinct from the two-cysteine-two-histidine zinc finger	170	5.5 Regulation of transcription by RNA polymerases I and III	199
The leucine zipper is a dimerization domain which allows DNA binding by the adjacent basic domain	175	Transcription by RNA polymerases I and III can be regulated by altering the expression or activity of components of their basal transcriptional complexes	199
In some transcription factors, the basic DNA-binding domain is found associated with a helix-loop-helix dimerization domain	176	Regulation of transcription by RNA polymerase III can involve specific transcription factors binding to RNA as well as to DNA	200
Dimerization between factors provides an additional level of regulation	177	Transcription by RNA polymerases I and III can be regulated by alterations in chromatin structure	201
Other domains can also mediate DNA binding	178	Conclusions	202
The DNA sequence bound by a particular transcription factor and the effect of such binding is affected by the presence or absence of other regulatory factors	180	Key concepts	202
5.2 Activation of transcription	181	Further reading	203
Activation domains can be identified by "domain-swap" experiments	181	6 Post-transcriptional Processes	205
		Introduction	205
		6.1 Capping	205
		The capping process modifies the 5' end of the RNA transcript	205

3.6 The cap enhances translation of the mRNA by the ribosome	206	Key concepts	231
6.2 Polyadenylation	208	Further reading	231
The polyadenylation process modifies the 3' end of the RNA transcript	208	7 Post-transcriptional Regulation	233
Polyadenylation enhances the stability of the mRNA	208	Introduction	233
6.3 RNA splicing	210	7.1 Alternative RNA splicing	234
RNA splicing removes intervening sequences and joins exons together	210	RNA splicing can be regulated	234
Specific RNAs and proteins catalyze the process of RNA splicing within the spliceosome	210	Alternative splicing represents a major regulatory process which supplements transcriptional control	234
6.4 Coupling of transcription and RNA processing within the nucleus	215	Alternative RNA splicing involves specific splicing factors that promote or inhibit the use of specific splice sites	235
Transcriptional initiation and elongation are coupled to post-transcriptional processes	215	Factors regulating alternative splicing have been identified by genetic and biochemical methods	237
Post-transcriptional processes can interact with one another	218	The processes of transcription and alternative splicing interact with one another	241
6.5 RNA transport	218	Alternative RNA splicing is a very widely used method of supplementing transcriptional control	243
RNA transport is coupled to other post-transcriptional processes	218	7.2 Regulation of polyadenylation	245
6.6 Translation	220	Both the amount and the site of polyadenylation can be regulated	245
Translation of the mRNA takes place on cytoplasmic ribosomes	220	Alternative polyadenylation can lead to alternative splicing	246
Translational initiation involves initiation factors binding to the cap	222	7.3 RNA editing	249
Translation involves base pairing of triplet codons in the mRNA with tRNA anticodons	222	Some cases of RNA editing involve a change from a C to a U residue	249
Translational elongation involves tRNAs located in two distinct sites within the ribosome	225	Many cases of RNA editing involve a change from A to an I residue	250
Translational termination occurs at specific stop codons	226	7.4 Regulation of RNA transport	252
6.7 RNA degradation	228	Specific proteins can regulate the transport of individual mRNAs from nucleus to cytoplasm	252
RNA degradation occurs in both the nucleus and the cytoplasm	228	RNA transport processes can also regulate the location of individual mRNAs within the cytoplasm	254
RNA degradation in the cytoplasm involves prior deadenylation and decapping of the mRNA	229	RNA localization can be linked to other stages of gene expression	255
Abnormal mRNAs are degraded in the cytoplasm by a nonsense-mediated RNA decay process	229	7.5 Regulation of RNA stability	256
Conclusions	231	Gene regulation can involve alterations in RNA stability	256

Specific sequences in the mRNA are involved in the regulation of its stability	257	Conclusions	280
RNA stability changes supplement transcriptional control in cases where a rapid response is required	258	Key concepts	281
7.6 Regulation of translation	260	Further reading	282
Translational control occurs in specific situations such as fertilization	260	8 Gene Control and Cellular Signaling Pathways	283
Whole genome methods can be used to study translational control	260	Introduction	283
Translational control can involve either modifications in the cellular translational apparatus or specific proteins which recognize sequences in the target RNA	262	Transcription factors can be regulated by controlling their synthesis or by controlling their activity	283
Translational control can be produced by modifications in the cellular translation apparatus	262	Multiple mechanisms regulate transcription factor activity	285
Translational control can be produced by proteins binding to the 5' untranslated region of the RNA itself	265	8.1 Regulation of transcription factor activity by ligands which enter the cell	286
Translational control can also be produced by proteins binding to the 3' untranslated region of the RNA itself	268	Transcription factors can be activated by direct binding of ligands which enter the cell	286
Translational control frequently occurs when a rapid response is required but also occurs for the genes encoding some transcription factors	270	Members of the nuclear receptor family of transcription factors are activated by binding of the appropriate ligand	286
7.7 Post-transcriptional inhibition of gene expression by regulatory RNAs	271	Following ligand-mediated activation, the glucocorticoid receptor can repress as well as activate gene transcription	289
Small RNAs can inhibit gene expression post-transcriptionally	271	The heat-shock factor is activated by stressful stimuli and induces the transcription of genes encoding protective proteins	291
Small RNAs can induce mRNA degradation	272	8.2 Regulation of transcription factor activity by phosphorylation induced by extracellular signaling molecules	292
Small RNAs can repress mRNA translation	274	Transcription factors can be phosphorylated by receptor-associated kinases	292
miRNAs regulate gene expression at multiple levels	276	Transcription factors can be phosphorylated by kinases activated by specific intracellular second messengers such as cyclic AMP	293
The production of mature miRNAs can be regulated post-transcriptionally as well as at transcription	278	Transcription factors can be phosphorylated by signaling cascades consisting of several protein kinases	295
Long noncoding RNAs can interact with small RNAs to regulate their activity	279	Transcription factor activity can be regulated by phosphorylation of an inhibitory protein: the NF κ B/I κ B system	296
Long noncoding RNAs can directly regulate gene expression post-transcriptionally	280		

8.3 Regulation of transcription factor activity by other post-translational modifications	298	Conclusions	311
8.4 Regulation of transcription factor activity by signals which regulate precursor processing	301	Key concepts	314
Transcription factors can be activated by cleavage of a precursor which contains an inhibitory region	301	Further reading	314
Transcription factors can be activated by cleavage of a membrane-bound precursor	301	9 Gene Control In Embryonic Development	317
Cleavage of a transcription factor can convert it from an activator to a repressor	302	Introduction	317
Cleavage of a lipid link can be used to activate a transcription factor	303	Regulation of mRNA translation occurs following fertilization	317
8.5. Regulation of histone modification and chromatin structure by cellular signaling pathways	304	Transcriptional control processes activate the embryonic genome	318
Cellular signaling pathways can induce the post-translational modification of histones	304	The Oct4 and Cdx2 transcription factors regulate the differentiation of inner cell mass and trophectoderm cells	320
Histone modifications link intermediary metabolism and chromatin structure/gene expression	304	9.1 Regulation of gene expression in pluripotent embryonic stem cells	321
8.6 Regulation of post-transcriptional processes by cellular signaling pathways	305	Embryonic stem cells can differentiate into a wide variety of cell types	321
The PI3-kinase/Akt system plays a key role in regulating gene expression in response to growth factors or insulin	305	Several transcription factors which are expressed specifically in ES cells can together reprogram differentiated cells to an ES cell-like phenotype	322
Akt regulates RNA splicing by phosphorylating splicing factors	306	ES cell-specific transcription factors can activate or repress the expression of their target genes	324
Akt can differentially regulate microRNA activity by phosphorylating the Argonaute 2 protein	307	ES cell-specific transcription factors regulate genes encoding chromatin-modifying enzymes and miRNAs	326
Akt regulates mRNA translation via the TOR kinase which phosphorylates proteins involved in translation	307	ES cells have an unusual pattern of histone methylation	328
Akt/TOR can also stimulate mRNA translation by enhancing the transcription of genes encoding RNAs and proteins involved in protein synthesis	308	The polycomb complex regulates histone methylation in ES cells	329
A variety of kinases inhibit translation by phosphorylating eIF2	309	Polycomb protein complexes regulate the expression of microRNA genes in ES cells	331
Individual kinases can produce multi-level regulation of gene expression	310	Chromatin structure in ES cells is regulated by multiple effects on histones	332
		ES cells show very specific patterns of chromosome looping which bring together transcriptionally active loci	332
		The regulation of chromatin structure in ES cells plays a critical role in maintaining their undifferentiated nature and in their ability to differentiate into a variety of cell types	333
		Gene expression in ES cells is regulated at the post-transcriptional level as well as at transcription	333

9.2 Role of gene regulation in the development of <i>Drosophila melanogaster</i>	337	Key concepts	352
A gradient in expression of the Bicoid transcription factor plays a key role in defining the anterior-posterior axis in the early <i>Drosophila</i> embryo	337	Further reading	352
Bicoid activates a cascade of genes encoding other transcription factors producing a segmented pattern of <i>Eve</i> gene expression	338	10 Control of Cell Type-Specific Gene Expression	355
The Bicoid system involves both transcriptional and post-transcriptional regulation	339	Introduction	355
Homeodomain transcription factors specify segment identity in the <i>Drosophila</i> embryo	340	10.1 Regulation of gene expression in skeletal muscle cells	357
Protein-protein interactions control the effect of homeodomain-containing transcription factors on gene expression	341	The MyoD protein can induce muscle cell differentiation	357
9.3 Role of homeodomain factors in mammalian development	341	MyoD is a basic helix-loop-helix transcription factor which is able to regulate gene expression	358
Homeodomain transcription factors are also found in mammals	341	MyoD is regulated by controlling both its synthesis and its activity	360
Mammalian <i>Hox</i> genes are expressed in specific regions of the developing embryo	342	Other muscle-specific transcription factors can induce muscle cell differentiation	361
Transcription of individual <i>Hox</i> genes is regulated by gene-specific regulatory regions	342	MEF2 proteins are downstream regulators of muscle-cell specific gene transcription	363
<i>Hox</i> gene transcription is also dependent on the position of the gene in the <i>Hox</i> gene cluster	344	10.2 Regulation of gene expression in neuronal cells	366
Changes in chromatin structure play a key role in activating the expression of <i>Hox</i> genes and in their spatial/temporal pattern of gene expression	344	Basic helix-loop-helix transcription factors are also involved in neuronal differentiation	366
<i>Hox</i> gene expression is also regulated at the post-transcriptional level by specific miRNAs	345	The REST transcription factor represses the expression of neuronal genes	369
Differential regulation of different <i>Hox</i> genes by Sonic Hedgehog controls the differentiation of cells in the neural tube	346	Neuronal cells express specific alternative splicing factors	371
Regulation of <i>Hox</i> gene expression by Sonic Hedgehog is also involved in limb formation	348	Translational control plays a key role in synaptic plasticity in neuronal cells	373
Conclusions	349	miRNAs play a key role in the regulation of neuronal gene expression	375
		10.3 Regulation of yeast mating type	377
		Yeast cells can be a or α in mating type	377
		Mating-type switching is controlled by regulating the transcription of the <i>HO</i> gene	378
		The SBF transcription factor activates <i>HO</i> transcription only in the G1 phase of the cell cycle	378
		The Ash-1 transcription factor represses <i>HO</i> transcription in daughter cells	379

The α and α gene products are homeodomain-containing transcription factors	379	11.3 Tumor suppressor transcription factors	402
The $\alpha 1$ and $\alpha 2$ proteins interact with the MCM1 transcription factor to respectively activate α -specific genes and repress α -specific genes	380	Tumor suppressor genes encode proteins which restrain cellular growth	402
The $\alpha 1$ factor plays a key role in repressing haploid-specific genes in diploid cells	381	The p53 protein is a DNA-binding transcription factor	402
The yeast mating-type system offers insights of relevance to multicellular organisms	383	The retinoblastoma protein interacts with other proteins to regulate transcription	407
Conclusions	385	Other tumor suppressor proteins also regulate transcription	409
Key concepts	386	11.4 The epigenome and cancer	412
Further reading	387	Oncogenic and tumor suppressor proteins can modulate chromatin structure	412
11 Gene Regulation and Cancer	389	Alterations in DNA methylation patterns play an important role in cancer	413
Introduction	389	Components of nucleosome remodeling complexes are mutated in different human cancers	414
11.1 Gene regulation and cancer	389	11.5 Post-transcriptional processes and cancer	414
Oncogenes were originally identified in cancer-causing viruses	389	Alterations in mRNA translation in cancer can be produced by enhancing the activity of RNA polymerases I and III	414
Cellular proto-oncogenes are present in the genome of normal cells	390	Alterations in mRNA translation in cancer can be produced directly by changes in ribosomal proteins	415
Cellular proto-oncogenes can cause cancer when they are overexpressed or mutated	391	Other post-transcriptional processes are also involved in cancer	416
Elevated expression of oncogenes can be produced by viruses or by cellular mechanisms	392	11.6 Regulatory RNAs and cancer	417
A variety of cellular mechanisms mediate enhanced expression of proto-oncogenes in different cancers	392	Different miRNAs can act to promote or inhibit cancer development	418
11.2 Oncogenic transcription factors	394	Different lncRNAs can both promote or inhibit cancer development	419
The Fos and Jun oncogene proteins are cellular transcription factors which can cause cancer when overexpressed	394	Conclusions	421
The v-erbA oncogene protein is a mutant form of the cellular thyroid hormone receptor	397	Key concepts	421
Other transcription factor-related oncogenes are overexpressed due to chromosomal translocations	399	Further reading	422
Chromosomal translocations can also produce novel oncogenic fusion proteins involving transcription factors	400	12 Gene Regulation and Human Disease	425
		Introduction	425
		12.1 Transcription and human disease	425
		12.2 The epigenome and human disease	430

12.3 Post-transcriptional processes and human disease	433	Transcription factors can repress gene expression as well as activating it	458
12.4 Regulatory RNAs and human disease	437	DNA-binding transcription factors interact with co-activators/co-repressors and with regulators of chromatin structure	458
A variety of mechanisms underlie the involvement of miRNAs in human diseases	437	Histone modifications play a central role in the regulation of chromatin structure	459
Long noncoding RNAs are involved in human disease at both the transcriptional and post-transcriptional levels	438	Co-activators/co-repressors link together different signaling pathways by interacting with multiple transcription factors	460
12.5 Infectious diseases and cellular gene expression	439	Gene regulation is highly complex and involves both transcriptional and post-transcriptional regulation	461
12.6 Gene regulation and therapy of human disease	442	Both small and large noncoding RNAs play a central role in regulating gene expression	461
Therapy could be achieved by altering the expression of transcription factors	442	Regulatory elements, chromatin structure, and DNA-binding proteins can be mapped across entire genomes	462
Therapy could be achieved by altering the activity of transcription factors	442	High-throughput methods are providing a full understanding of the various stages of gene expression and its regulation	463
Therapy could be achieved by targeting proteins which alter chromatin structure	444	Gene control is necessarily a highly complex process	463
Therapy could be achieved using designer zinc fingers to alter gene transcription	448	Further reading	465
Therapy could be achieved by modulating RNA splicing	449	Glossary	467
Therapy could be achieved by targeting regulatory RNAs	451	Index	489
Conclusions	453		
Key concepts	454		
Further reading	455		
13 Conclusions and Future Prospects	457		
Conclusions and future prospects	457		
Transcription factors interact with one another to regulate transcription	457		