

# Brief Contents

|                                                                                 |            |
|---------------------------------------------------------------------------------|------------|
| <b>Chapter 1: Introduction to Cell Signaling</b>                                | <b>1</b>   |
| <b>Chapter 2: Principles and Mechanisms of Protein Interactions</b>             | <b>21</b>  |
| <b>Chapter 3: Signaling Enzymes and Their Allosteric Regulation</b>             | <b>43</b>  |
| <b>Chapter 4: Role of Post-Translational Modifications in Signaling</b>         | <b>85</b>  |
| <b>Chapter 5: Subcellular Localization of Signaling Molecules</b>               | <b>115</b> |
| <b>Chapter 6: Second Messengers: Small Signaling Mediators</b>                  | <b>135</b> |
| <b>Chapter 7: Membranes, Lipids, and Enzymes That Modify Them</b>               | <b>155</b> |
| <b>Chapter 8: Information Transfer Across the Membrane</b>                      | <b>177</b> |
| <b>Chapter 9: Regulated Protein Degradation</b>                                 | <b>217</b> |
| <b>Chapter 10: The Modular Architecture and Evolution of Signaling Proteins</b> | <b>243</b> |
| <b>Chapter 11: Information Processing by Signaling Devices and Networks</b>     | <b>275</b> |
| <b>Chapter 12: How Cells Make Decisions</b>                                     | <b>305</b> |
| <b>Chapter 13: Methods for Studying Signaling Proteins and Networks</b>         | <b>345</b> |
| <b>Glossary</b>                                                                 | <b>375</b> |
| <b>Index</b>                                                                    | <b>385</b> |

# Detailed Contents

## Chapter 1

### Introduction to Cell Signaling

#### WHAT IS CELL SIGNALING?

- All cells have the ability to respond to their environment 2
- Cells must perceive and respond to a wide range of signals 3
- Signaling systems need to solve a number of common problems 4

#### THE FUNDAMENTAL ROLE OF SIGNALING IN BIOLOGICAL PROCESSES

- Work in many different fields converged to reveal the underlying mechanisms of signaling 6
- Despite the diversity of signaling pathways and mechanisms, fundamental commonalities have emerged 7
- Signaling must operate at multiple scales in space and time 9

#### THE MOLECULAR CURRENCIES OF INFORMATION PROCESSING

- Information is transferred by changes in the state of proteins 11
- There is a limited number of ways in which the state of proteins can change 12
- Most changes in state involve simultaneous changes in several different currencies 14

#### LINKING SIGNALING NODES INTO PATHWAYS AND NETWORKS

- Information transfer involves linking different changes of state together 15
- Multiple state changes are linked together to generate pathways and networks 16
- Cellular information-processing systems have a hierarchical architecture 17
- Summary 18
- Questions 18
- References 19

## Chapter 2

### Principles and Mechanisms of Protein Interactions

#### PROPERTIES OF PROTEIN-PROTEIN INTERACTIONS

- Changes in protein binding have both direct and indirect functional consequences 22
- Protein binding can be mediated by broad interaction surfaces or by short, linear peptides 23

- The affinity and specificity of an interaction determine how likely it is to occur in the cell 24
- The strength of a binding interaction is defined by the dissociation constant ( $K_d$ ) 25
- The dissociation constant is related to the binding energy of the interaction 27
- The dissociation constant is also related to rates of binding and dissociation 28

#### PROTEIN INTERACTIONS IN THEIR CELLULAR AND MOLECULAR CONTEXT

- The apparent dissociation constant can be strongly affected by the local cellular environment and other binding partners 30
- Ideal affinity and specificity depends on biological function and ligand concentrations 31
- There are functional constraints on interaction affinities and specificities 32
- Interaction affinity and specificity can be independently modulated 34
- Cooperativity involves the coupled binding of multiple ligands 35
- Diverse molecular mechanisms underlie cooperativity 36
- Cooperative binding has a variety of functional consequences 36
- Protein assemblies differ in their stability and homogeneity 37
- Summary 38
- Questions 38
- References 40

## Chapter 3

### Signaling Enzymes and Their Allosteric Regulation

#### PRINCIPLES OF ENZYME CATALYSIS

- Enzymes have a number of properties that make them useful for transmitting signals in the cell 44
- Enzymes use a variety of mechanisms to enhance the rate of chemical reactions 45
- Enzymes can drive reactions in one direction by energetic coupling 46

#### ALLOSTERIC CONFORMATIONAL CHANGES

- Conformational flexibility of proteins enables allosteric control 47

|                                                                                                                                               |    |                                                                                                         |           |
|-----------------------------------------------------------------------------------------------------------------------------------------------|----|---------------------------------------------------------------------------------------------------------|-----------|
| Signaling proteins employ diverse classes of conformational rearrangements                                                                    | 48 | Questions                                                                                               | 80        |
| PROTEIN PHOSPHORYLATION AS A REGULATORY MECHANISM                                                                                             | 49 | References                                                                                              | 82        |
| Phosphorylation can act as a regulatory mark                                                                                                  | 49 | <b>Chapter 4</b>                                                                                        |           |
| Phosphorylation can either disrupt or induce protein structure                                                                                | 50 | <b>Role of Post-Translational Modifications in Signaling</b>                                            | <b>85</b> |
| PROTEIN KINASES                                                                                                                               | 52 | THE LOGIC OF POST-TRANSLATIONAL REGULATION                                                              | 85        |
| The structure and catalytic mechanism of protein kinases are conserved                                                                        | 52 | Proteins can be covalently modified by the addition of simple functional groups                         | 86        |
| The activation loop and C-helix are conserved molecular levers that conformationally control kinase activity                                  | 54 | Proteins can also be covalently modified by the addition of sugars, lipids, and even proteins           | 87        |
| Insulin receptor kinase activity is controlled via activation-loop phosphorylation                                                            | 54 | Post-translational modifications can alter protein structure, localization, and stability               | 88        |
| Phosphorylation mediates long-range conformational regulation of Src family kinases                                                           | 55 | Post-translational control machinery often works as part of "writer/eraser/reader" systems              | 90        |
| Multiple binding interactions regulate protein kinase substrate specificity                                                                   | 56 | Post-translational modifications allow very rapid signaling and transmission of spatial information     | 92        |
| Protein kinases can be divided into nine families                                                                                             | 58 | INTERPLAY BETWEEN POST-TRANSLATIONAL MODIFICATIONS                                                      | 92        |
| PROTEIN PHOSPHATASES                                                                                                                          | 60 | A post-translational modification can promote or antagonize other modifications                         | 93        |
| Serine/threonine phosphatases are metalloenzymes                                                                                              | 60 | p53 is tightly regulated by a wide variety of post-translational modifications                          | 95        |
| Most tyrosine phosphatases utilize a catalytic cysteine residue                                                                               | 62 | The level and activity of p53 are regulated by ubiquitylation and acetylation                           | 96        |
| Tyrosine phosphatases are regulated by modular domains while serine/threonine phosphatases often associate with regulatory accessory subunits | 64 | Additional modifications further fine-tune p53 activity                                                 | 96        |
| G PROTEIN SIGNALING                                                                                                                           | 65 | PROTEIN PHOSPHORYLATION                                                                                 | 97        |
| G proteins are conformational switches controlled by two opposing enzymes                                                                     | 65 | Phosphorylation is often coupled with protein interactions                                              | 97        |
| The presence of the GTP $\gamma$ -phosphate determines the structure of G protein switch I and II regions                                     | 66 | Kinases and phosphatases vary in their substrate specificity                                            | 99        |
| There are two major classes of signaling G proteins                                                                                           | 67 | Multiple phosphorylation of proteins can arise by different mechanisms                                  | 100       |
| Subfamilies of small G proteins regulate diverse biological functions                                                                         | 67 | Histidine and other amino acids can be phosphorylated, especially in prokaryotes                        | 101       |
| Many upstream receptors feed into a small set of common heterotrimeric G proteins                                                             | 68 | Two-component systems and histidine phosphorylation are also present in eukaryotes                      | 103       |
| REGULATORY ENZYMES FOR G PROTEIN SIGNALING                                                                                                    | 70 | ADDITION OF UBIQUITIN AND RELATED PROTEINS                                                              | 104       |
| G-protein-coupled receptors act as GEFs for heterotrimeric G proteins                                                                         | 71 | Specialized enzymes mediate the addition and removal of ubiquitin                                       | 104       |
| Distinct GEF and GAP domains regulate specific small G protein families                                                                       | 71 | E3 ubiquitin ligases determine which proteins will be ubiquitylated                                     | 105       |
| GEFs catalyze GDP/GTP exchange by deforming the nucleotide-binding pocket                                                                     | 73 | Ubiquitin-binding domains read ubiquitin-mediated signals in diverse cellular activities                | 106       |
| GAPs order the catalytic machinery for hydrolysis                                                                                             | 74 | HISTONE ACETYLATION AND METHYLATION                                                                     | 107       |
| Regulators of G protein signaling (RGS) proteins act as GAPs for heterotrimeric G proteins                                                    | 75 | Chromatin structure is regulated by post-translational modification of histones and associated proteins | 108       |
| Additional mechanisms are used to fine-tune the activity of G proteins                                                                        | 75 | Two writer/eraser/reader systems are based on protein methylation and acetylation                       | 109       |
| SIGNALING ENZYME CASCADES                                                                                                                     | 75 | Chromatin modification in transcription is dynamic and leads to highly cooperative interactions         | 110       |
| The three-tiered MAP kinase cascade forms a signaling module in all eukaryotes                                                                | 76 | Summary                                                                                                 | 112       |
| Scaffold proteins often organize MAPK cascades                                                                                                | 77 | Questions                                                                                               | 113       |
| G protein activity can also be regulated by signaling cascades                                                                                | 79 | References                                                                                              | 114       |
| Summary                                                                                                                                       | 80 |                                                                                                         |           |

|                                                                                                          |            |
|----------------------------------------------------------------------------------------------------------|------------|
| <b>Chapter 5</b>                                                                                         |            |
| <b>Subcellular Localization of Signaling Molecules</b>                                                   | <b>115</b> |
| LOCALIZATION AS A SIGNALING CURRENCY                                                                     | 115        |
| Changes in subcellular localization can transmit information                                             | 116        |
| Subcellular localization can be regulated by a variety of mechanisms                                     | 117        |
| CONTROL OF NUCLEAR LOCALIZATION                                                                          | 117        |
| Short, modular peptide motifs direct nuclear import and export                                           | 118        |
| Nuclear transport is controlled by shuttle proteins and the G protein Ran                                | 118        |
| Phosphorylation of transcription factor Pho4 regulates nuclear import and export                         | 119        |
| Nuclear import of STATs is regulated by phosphorylation and conformational change                        | 120        |
| Localization of MAP kinases is regulated by association with nuclear and cytosolic binding partners      | 121        |
| Notch nuclear localization is regulated by proteolytic cleavage                                          | 122        |
| CONTROL OF MEMBRANE LOCALIZATION                                                                         | 122        |
| Proteins can span the membrane or be associated with it peripherally                                     | 122        |
| Proteins can be covalently modified with lipids after translation                                        | 123        |
| Modular lipid-binding domains are important for regulated association of proteins with membranes         | 124        |
| Some lipid-modified proteins can reversibly associate with membranes                                     | 125        |
| Coupling effector protein activation to membrane recruitment is a common theme in signaling              | 126        |
| Akt kinase is regulated by membrane recruitment and phosphorylation                                      | 126        |
| MODULATION OF SIGNALING BY MEMBRANE TRAFFICKING                                                          | 127        |
| Proteins can be internalized by a variety of mechanisms                                                  | 127        |
| Internalization of receptors can modulate signal transduction                                            | 128        |
| TGF $\beta$ signaling output depends on the mechanism of receptor internalization                        | 129        |
| Retrograde signaling allows effects distant from the site of ligand binding                              | 130        |
| Ras isoforms in distinct subcellular locations have different signaling outputs                          | 130        |
| Summary                                                                                                  | 132        |
| Questions                                                                                                | 132        |
| References                                                                                               | 132        |
| <b>Chapter 6</b>                                                                                         |            |
| <b>Second Messengers: Small Signaling Mediators</b>                                                      | <b>135</b> |
| PROPERTIES OF SMALL SIGNALING MEDIATORS                                                                  | 135        |
| Small signaling mediators are controlled by an interplay of their production and elimination             | 136        |
| Small signaling mediators exert their effects by binding downstream effectors                            | 136        |
| Small signaling mediators can lead to fast, distant, and amplified signal transmission                   | 137        |
| Small signaling mediators can generate complex temporal and spatial patterns                             | 138        |
| CLASSES OF SMALL SIGNALING MEDIATORS                                                                     | 139        |
| Small signaling mediators have a wide range of physical properties                                       | 140        |
| The cyclic nucleotides cAMP and cGMP are produced by cyclase enzymes and destroyed by phosphodiesterases | 140        |
| Cyclic nucleotides regulate diverse cellular activities                                                  | 141        |
| The regulatory (R) subunit of protein kinase A is a conformational sensor of cAMP binding                | 142        |
| Some small signaling mediators are derived from membrane lipids                                          | 143        |
| PLC generates two signaling mediators, IP <sub>3</sub> and DAG                                           | 144        |
| Activation of protein kinase C is regulated by IP <sub>3</sub> and DAG                                   | 144        |
| CALCIUM SIGNALING                                                                                        | 145        |
| Activation of Ca <sup>2+</sup> channels is a common means of regulation                                  | 146        |
| Ca <sup>2+</sup> influx is rapid and local                                                               | 147        |
| Calmodulin is a conformational sensor of intracellular calcium levels                                    | 147        |
| Signaling can lead to propagating Ca <sup>2+</sup> waves                                                 | 148        |
| SPECIFICITY AND REGULATION                                                                               | 149        |
| Scaffold proteins can increase input and output specificity of small-molecule signaling                  | 150        |
| AKAP scaffold proteins can also regulate dynamics of cAMP signaling                                      | 150        |
| Summary                                                                                                  | 152        |
| Questions                                                                                                | 152        |
| References                                                                                               | 153        |
| <b>Chapter 7</b>                                                                                         |            |
| <b>Membranes, Lipids, and Enzymes That Modify Them</b>                                                   | <b>155</b> |
| BIOLOGICAL MEMBRANES AND THEIR PROPERTIES                                                                | 155        |
| Biological membranes consist of a variety of polar lipids                                                | 156        |
| Structural properties of membrane lipids favor the formation of bilayers                                 | 157        |
| The composition of the membrane determines its physical properties                                       | 158        |
| There are fundamental differences between biochemistry in solution and on the membrane                   | 160        |
| LIPID-MODIFYING ENZYMES USED IN SIGNALING                                                                | 161        |
| Cleavage of membrane lipids by phospholipases generates a variety of bioactive products                  | 161        |
| A variety of lipid kinases and phosphatases are involved in signaling                                    | 163        |

|                                                                                                               |     |                                                                                                                    |     |
|---------------------------------------------------------------------------------------------------------------|-----|--------------------------------------------------------------------------------------------------------------------|-----|
| <b>EXAMPLES OF MAJOR LIPID SIGNALING PATHWAYS</b>                                                             | 164 | The voltage-gated potassium channel provides clues to mechanisms of gating and ion specificity                     | 200 |
| Phosphoinositides can serve as membrane binding sites and as a source of signaling mediators                  | 164 | Ligand-gated ion channels play a central role in neurotransmission                                                 | 202 |
| Phosphoinositide species provide a set of membrane binding signals                                            | 166 | <b>MEMBRANE-PERMEABLE SIGNALING</b>                                                                                | 204 |
| Phospholipase D generates the important signaling mediator, phosphatidic acid (PA)                            | 168 | Nitric oxide mediates short-range signaling in the vascular system                                                 | 204 |
| Phospholipase D plays a role in mTOR signaling                                                                | 169 | O <sub>2</sub> binding regulates the response to hypoxia                                                           | 205 |
| The metabolism of sphingomyelin generates a host of signaling mediators                                       | 170 | The receptors for steroid hormones are transcription factors                                                       | 206 |
| Phospholipase A <sub>2</sub> generates the precursor for a family of potent inflammatory mediators            | 172 | <b>DOWN-REGULATION OF RECEPTOR SIGNALING</b>                                                                       | 208 |
| Summary                                                                                                       | 174 | Ubiquitylation regulates the endocytosis, recycling, and degradation of cell-surface receptors                     | 209 |
| Questions                                                                                                     | 174 | G protein coupled receptors are desensitized by phosphorylation and adaptor binding                                | 211 |
| References                                                                                                    | 174 | Summary                                                                                                            | 213 |
| <b>Chapter 8</b>                                                                                              |     | Questions                                                                                                          | 213 |
| <b>Information Transfer Across the Membrane</b>                                                               | 177 | References                                                                                                         | 215 |
| <b>PRINCIPLES OF TRANSMEMBRANE SIGNALING</b>                                                                  | 177 | <b>Chapter 9</b>                                                                                                   |     |
| The cell must process and respond to a diversity of environmental cues                                        | 178 | <b>Regulated Protein Degradation</b>                                                                               | 217 |
| Three general strategies are used to transfer information across the membrane                                 | 179 | <b>GENERAL PROPERTIES AND EXAMPLES OF SIGNAL-REGULATED PROTEOLYSIS</b>                                             | 217 |
| Many drugs target receptors                                                                                   | 180 | Proteases are a diverse group of enzymes                                                                           | 218 |
| <b>TRANSDUCTION STRATEGIES USED BY TRANSMEMBRANE RECEPTORS</b>                                                | 180 | Blood coagulation is regulated by a cascade of proteases                                                           | 219 |
| Receptors with multiple membrane-spanning segments undergo conformational changes upon ligand binding         | 180 | Regulated proteolysis by metalloproteases can generate signaling molecules and alter the extracellular environment | 220 |
| Receptors with a single membrane-spanning segment form higher-order assemblies upon ligand binding            | 181 | ADAMs regulate signaling pathways by cleaving membrane-associated proteins                                         | 221 |
| Receptor clustering confers advantages for signal propagation                                                 | 182 | MMPs participate in remodeling the extracellular environment                                                       | 222 |
| <b>G-PROTEIN-COUPLED RECEPTORS</b>                                                                            | 184 | Proteolysis activates the thrombin receptor                                                                        | 223 |
| G-protein-coupled receptors have intrinsic enzymatic activity                                                 | 184 | Regulated intramembrane proteolysis (RIP) is an essential step in signaling by some receptors                      | 224 |
| Signaling by GPCRs can be very fast and lead to enormous signal amplification                                 | 186 | <b>UBIQUITIN AND THE PROTEASOME DEGRADATION PATHWAY</b>                                                            | 225 |
| <b>TRANSMEMBRANE RECEPTORS ASSOCIATED WITH ENZYMATIC ACTIVITY</b>                                             | 186 | The proteasome is a specialized molecular machine that degrades intracellular proteins                             | 225 |
| Receptor tyrosine kinases control important cell fate decisions in multicellular eukaryotes                   | 186 | The cell cycle is controlled by two large ubiquitin-conjugating complexes                                          | 226 |
| TGF $\beta$ receptors are serine/threonine kinases that activate transcription factors                        | 187 | SCF recognizes specific phosphorylated proteins, targeting them for destruction                                    | 227 |
| Some receptors have intrinsic protein phosphatase or guanylyl cyclase activity                                | 189 | Two APC species act at distinct points in the cell cycle                                                           | 228 |
| Noncovalent coupling of receptors to protein kinases is a common signaling strategy                           | 189 | NF- $\kappa$ B is controlled by regulated degradation of its inhibitor                                             | 230 |
| Some receptors use complex activation pathways that involve both kinase activation and proteolytic processing | 193 | <b>CASPASE-MEDIATED CELL DEATH PATHWAYS</b>                                                                        | 232 |
| Wnt and Hedgehog are two important signaling pathways in development                                          | 194 | Apoptosis is an orderly and highly regulated form of cell death                                                    | 232 |
| A variety of receptors couple to proteolytic activities                                                       | 197 | The activity of caspases is tightly regulated                                                                      | 233 |
| <b>GATED CHANNELS</b>                                                                                         | 199 | The extrinsic pathway links cell death receptors to caspase activation                                             | 235 |
| Gated channels share a similar overall structure                                                              | 199 | Mitochondria orchestrate the intrinsic cell death pathway                                                          | 238 |
|                                                                                                               |     | Summary                                                                                                            | 241 |

|                                                                                                                  |            |                                                                                     |            |
|------------------------------------------------------------------------------------------------------------------|------------|-------------------------------------------------------------------------------------|------------|
| Questions                                                                                                        | 241        | Certain plant protein kinases are regulated by modular light-gated domains          | 267        |
| References                                                                                                       | 242        | Regulation of the neutrophil NADPH oxidase by modular interactions                  | 268        |
| <b>Chapter 10</b>                                                                                                |            | <b>CREATING NEW FUNCTIONS THROUGH DOMAIN RECOMBINATION</b>                          | 269        |
| <b>The Modular Architecture and Evolution of Signaling Proteins</b>                                              | <b>243</b> | Some modular domain rearrangements can lead to cancer                               | 269        |
| MODULAR PROTEIN DOMAINS                                                                                          | 244        | Modules can be recombined experimentally to engineer new signaling behaviors        | 270        |
| Protein domains usually have a globular structure                                                                | 244        | Summary                                                                             | 272        |
| Bioinformatic approaches can identify protein domains                                                            | 244        | Questions                                                                           | 272        |
| Domains can be composed of several smaller repeats                                                               | 245        | References                                                                          | 273        |
| Protein domains often act as recognition modules                                                                 | 246        | <b>Chapter 11</b>                                                                   |            |
| INTERACTION DOMAINS THAT RECOGNIZE POST-TRANSLATIONAL MODIFICATIONS                                              | 249        | <b>Information Processing by Signaling Devices and Networks</b>                     | <b>275</b> |
| SH2 domains bind phosphotyrosine-containing sites                                                                | 249        | SIGNALING SYSTEMS AS INFORMATION-PROCESSING DEVICES                                 | 276        |
| Some SH2 domains are elements of larger binding structures                                                       | 252        | Signaling devices can be considered as state machines                               | 276        |
| Several different types of interaction domains recognize phosphotyrosine                                         | 252        | Signaling devices are organized in a hierarchical fashion                           | 277        |
| Multiple domains recognize motifs phosphorylated on serine/threonine                                             | 254        | Signaling devices face a variety of challenges in input detection                   | 278        |
| 14-3-3 proteins recognize specific phosphoserine/phosphothreonine motifs                                         | 254        | Proteins can function as simple signaling devices                                   | 279        |
| Interaction domains recognize acetylated and methylated sites                                                    | 255        | INTEGRATING MULTIPLE SIGNALING INPUTS                                               | 281        |
| Ubiquitylation regulates protein-protein interactions                                                            | 256        | Logic gates process information from multiple inputs                                | 281        |
| INTERACTION DOMAINS THAT RECOGNIZE UNMODIFIED PEPTIDE MOTIFS OR PROTEINS                                         | 257        | Simple peptide motifs can integrate multiple post-translational modification inputs | 282        |
| Proline-rich sequences are favorable recognition motifs                                                          | 257        | Cyclin-dependent kinase is an allosteric signal-integrating device                  | 283        |
| SH3 domains bind proline-rich motifs                                                                             | 258        | Modular signaling proteins can integrate multiple inputs                            | 284        |
| PDZ domains recognize C-terminal peptide motifs                                                                  | 258        | Transcriptional promoters can integrate input from multiple signaling pathways      | 285        |
| Protein interaction domains can form dimers or oligomers                                                         | 259        | RESPONDING TO THE STRENGTH OR DURATION OF AN INPUT                                  | 286        |
| INTERACTION DOMAINS THAT RECOGNIZE PHOSPHOLIPIDS                                                                 | 260        | Signaling systems can respond to signal amplitude in a graded or a digital manner   | 288        |
| PH domains form a major class of phosphoinositide-binding domains                                                | 260        | An enzyme can behave as a switch through cooperativity                              | 289        |
| FYVE domains are phospholipid-binding domains found in endocytic proteins                                        | 261        | Networks can also yield switchlike activation                                       | 290        |
| BAR domains bind and stabilize curved membranes                                                                  | 262        | Signaling systems can distinguish between transient and sustained input             | 292        |
| CREATING COMPLEX FUNCTIONS BY COMBINING INTERACTION DOMAINS                                                      | 262        | MODIFYING THE STRENGTH OR DURATION OF OUTPUT                                        | 294        |
| Recombination of domains occurs through evolution                                                                | 262        | Signaling pathways often amplify signals as they are transmitted                    | 294        |
| Combinations of interaction domains or motifs can be used as a scaffold for the assembly of signaling complexes  | 263        | Negative feedback allows fine-tuning of output                                      | 294        |
| Scaffold proteins containing PDZ domains organize cell-cell signaling complexes such as the postsynaptic density | 264        | Adaptation allows cells to control output duration                                  | 296        |
| Proteins with multiple phosphotyrosine motifs function as dynamically regulated scaffolds                        | 265        | Feedback can cause output levels to oscillate between two stable states             | 299        |
| RECOMBINING INTERACTION AND CATALYTIC DOMAINS TO BUILD COMPLEX ALLOSTERIC SWITCH PROTEINS                        | 266        | Bistable responses also underlie more permanent outputs                             | 301        |
| Many signaling enzymes are allosteric switches                                                                   | 266        | Summary                                                                             | 303        |
| 14-3-3 Protein regulates the Raf kinase by coordinately binding two phosphorylation sites                        | 267        | Questions                                                                           | 303        |
|                                                                                                                  |            | References                                                                          | 303        |

|                                                                                                                           |            |                                                                                                         |            |
|---------------------------------------------------------------------------------------------------------------------------|------------|---------------------------------------------------------------------------------------------------------|------------|
| <b>Chapter 12</b>                                                                                                         |            |                                                                                                         |            |
| <b>How Cells Make Decisions</b>                                                                                           | <b>305</b> | <b>Chapter 13</b>                                                                                       |            |
| VERTEBRATE VISION                                                                                                         | 307        | <b>Methods for Studying Signaling Proteins and Networks</b>                                             | <b>345</b> |
| Organ: vertebrate eye                                                                                                     | 308        | BIOCHEMICAL AND BIOPHYSICAL ANALYSIS OF PROTEINS                                                        | 345        |
| Cell: photoreceptor cell                                                                                                  | 308        | Analytical methods can determine quantitative binding parameters                                        | 345        |
| Molecular Network: visual transduction cascade                                                                            | 309        | Michaelis–Menten analysis provides a way to measure the catalytic power of enzymes                      | 347        |
| How does the photoreceptor cell detect light and convert it to a biochemical signal?                                      | 310        | Methods to determine and analyze protein conformation are central to the study of signaling             | 349        |
| How is the photoreceptor cell able to detect low light, even a single photon?                                             | 311        | X-ray crystallography provides high-resolution protein structures                                       | 352        |
| How can the response be so rapid?                                                                                         | 312        | Nuclear magnetic resonance (NMR) can reveal the dynamic structure of small proteins                     | 353        |
| How does the photoreceptor cell reset itself to enable detection of further changes in light?                             | 312        | Electron microscopy can map the shape of very large protein complexes                                   | 353        |
| Summary                                                                                                                   | 314        | Specialized spectroscopic methods can be used to study protein dynamics                                 | 354        |
| References                                                                                                                | 314        | MAPPING PROTEIN INTERACTIONS AND LOCALIZATION                                                           | 355        |
| PDGF SIGNALING                                                                                                            | 315        | Interacting proteins can be identified by isolating protein complexes from cell extracts                | 355        |
| Tissue: process of wound healing                                                                                          | 316        | Binding partners can be identified by screening large libraries of genes                                | 356        |
| Cell: fibroblast response to wounding & platelet activation                                                               | 316        | Direct protein–protein interactions can be detected by solid-phase screening                            | 357        |
| Molecular Network: control of fibroblast proliferation                                                                    | 317        | Fluorescent protein tags are used to locate and track proteins in living cells                          | 357        |
| How do fibroblasts detect the local occurrence of a wound?                                                                | 318        | Protein–protein interactions can be visualized directly in living cells                                 | 359        |
| How are PDGF signals propagated within the cell to generate outputs such as cell proliferation?                           | 319        | METHODS TO PERTURB CELL SIGNALING NETWORKS AND MONITOR CELLULAR RESPONSES                               | 360        |
| How is misactivation of the proliferation response prevented?                                                             | 320        | Genetic and pharmacological methods can be used to perturb networks                                     | 361        |
| How is the proliferative response terminated?                                                                             | 321        | Chemical dimerizers and optogenetic proteins provide a dynamic way to artificially activate pathways    | 362        |
| Summary                                                                                                                   | 322        | cDNA microarrays and high-throughput sequencing are used to monitor the transcriptional state of a cell | 363        |
| References                                                                                                                | 322        | Modification-specific antibodies provide a method to track post-translational changes                   | 364        |
| THE CELL CYCLE                                                                                                            | 323        | Mass spectrometry is the workhorse for identification of proteins and their modifications               | 366        |
| Cell: distinct phases of the cell cycle                                                                                   | 324        | Live-cell time-lapse microscopy provides a way to track the dynamics of single-cell responses           | 368        |
| Molecular Network: cyclin-dependent kinase (Cdk) is a central switch whose activity is modulated by the different cyclins | 325        | Biosensors allow signaling activity to be monitored in living cells                                     | 369        |
| How are sharp and committed transitions between cell cycle phases achieved?                                               | 326        | Flow cytometry provides a method to analyze rapidly single-cell responses                               | 371        |
| How does the cell cycle ensure that each transition proceeds only under appropriate conditions?                           | 329        | Questions                                                                                               | 372        |
| Summary                                                                                                                   | 331        | References                                                                                              | 373        |
| References                                                                                                                | 331        | <b>Glossary</b>                                                                                         | <b>375</b> |
| T CELL SIGNALING                                                                                                          | 333        | <b>Index</b>                                                                                            | <b>385</b> |
| Organism: launching the adaptive immune response                                                                          | 334        |                                                                                                         |            |
| Cell: engagement of T cell and antigen presenting cell                                                                    | 335        |                                                                                                         |            |
| Molecular Network: T cell receptor (TCR) signaling network                                                                | 336        |                                                                                                         |            |
| How does the T cell receptor transmit signals after peptide/MHC recognition?                                              | 338        |                                                                                                         |            |
| How does the T cell prevent misactivation?                                                                                | 339        |                                                                                                         |            |
| How does the T cell launch a robust response when stimulated by as few as ten antigenic peptide complexes?                | 340        |                                                                                                         |            |
| How might the T cell discriminate between antigenic and non-antigenic peptides?                                           | 342        |                                                                                                         |            |
| Summary                                                                                                                   | 344        |                                                                                                         |            |
| References                                                                                                                | 344        |                                                                                                         |            |