

SYNOPTIC INDEX

*The numbers in the first column refer to pages in this book,
those in the second column refer to pages in Engler's Syllabus
(Ed. 7, 1912).*

Division 1. EMBRYOPHYTA ASIPHONOGAMA (ARCHEGONIATAE)	1	74
[Subdivision 1. Bryophyta]	1	75
Subdivision 2. PTERIDOPHYTA	1	88
Class 1. EQUISETARIAE (Equisetales in Syllabus)	1	97
Order Equisetales (Euequisetales in Syllabus)	2	97
Fam. EQUISETACEAE	2	97
Class 2. LYCOPODIARIAE (Lycopodiales in Syllabus)	2	98
Order 1. Lycopodiales (Lycopodiales eligulatae and ligulatae of Syllabus)	2	98-99
Fam. 1. LYCOPODIACEAE	2	98
Fam. 2. SELAGINELLACEAE	3	99
Order 2. Isoëtales (Class in Syllabus)	3	101
Fam. ISOËTACEAE	3	101
Class 3. FILICARIAE (Filicales in Syllabus).	3	89

Synoptic Index

Order 1. Filicales (Filicales Leptosporangiatæ in Syllabus)	4	89
Fam. 1. OSMUNDACEAE	4	94
Fam. 2. HYMENOPHYLLACEAE	5	90
Fam. 3. POLYPODIACEAE	5	90
Fam. 4. MARSILIACEAE	7	95
Fam. 5. SALVINIACEAE	7	95
Order 2. Ophioglossales	7	96
Fam. OPHIOGLOSSACEAE	8	97
Division 2. EMBRYOPHYTA SIPHONOGAMA (<i>Phanerogamæ</i> auct.)	8	102
Subdivision 1. GYMNOSPERMÆ	8	103
Class 1. CONIFERÆ	9	106
Fam. 1. TAXACEAE	9	106
Fam. 2. PINACEAE	9	108
Subdivision 2. ANGIOSPERMÆ	10	113
Class 1. MONOCOTYLEDONES	10	117
A. Orders with preponderating inconstancy in the number of floral leaves	10	117
a. Typical achlamydeous (not apopetalous) flowers still occur	10	117
a. Achlamydeous flowers predominate. Great inconstancy in number of stamens and also of carpels	10	117
Order 1. Pandanales	10	117
Fam. 1. TYPHACEAE	11	118
Fam. 2. SPARGANIACEAE	11	118
β. Naked flowers still present: but in this order there occur all possible gradations from Achlamydy to Heterochlamydy, also from Hypogyny to Epigyny. Sporophylls definite or indefinite	11	118

Synoptic Index

xi

Order 2. Alismatales	11	118
Fam. 1. POTAMOGETONACEAE	11	119
Fam. 2. NAIADACEAE	12	120
Fam. 3. *APONOGETONACEAE	12	120
Fam. 4. SCHEUCHZERIAACEAE	13	120
Fam. 5. ALISMATAACEAE	13	120
Fam. 6. BUTOMACEAE	14	121
Fam. 7. HYDROCHARITACEAE	14	121
γ. <i>Naked flowers dominant. Stamens rarely, and carpels never, indefinite</i>	15	123
Order 3. Glumiflorae	15	123
Fam. 1. GRAMINACEAE	15	123
Fam. 2. CYPERACEAE	20	127
b. <i>Achlamydeous flowers rare: when they occur they are mostly due to reduction and are correlated with spathe-development. Definite number of stamens and carpels dominant, but numerous stamens and more than three carpels often occur</i>	21	128
Order 4. Spathiflorae	22	133
Fam. 1. ARACEAE	22	133
Fam. 2. LEMNACEAE	23	136
B. Orders with typically pentacyclic flowers. Whorls typically isomerous, mostly 3-merous	23	137
a. <i>Flowers homochlamydeous to heterochlamydeous, very rarely by reduction naked. Bracteoid perianths still occur but petaloid perianths predominate. Hypogyny and Actinomorphy preponderate</i>	23	137
Order 5. Farinosae	23	137
Fam. ERIOCAULACEAE	23	138

Synoptic Index

Order 6. Liliiflorae	24	141
Fam. 1. JUNCACEAE	24	141
Fam. 2. LILIACEAE	24	141
Fam. 3. AMARYLLIDACEAE	27	146
Fam. 4. DIOSCOREACEAE	28	147
Fam. 5. IRIDACEAE	28	148
<i>b. Flowers homochlamydeous to heterochlamydeous but the perianth is always petaloid. Epigyny throughout. Zygomorphy dominant</i>		
Order 7. Microspermae	29	151
Fam. ORCHIDACEAE	30	152
Class 2. DICOTYLEDONES	33	156
Subclass 1. ARCHICHLAMYDEAE	33	156
<i>a. Amentiflorae (Moss). Mostly trees or shrubs. ♂ flowers usually, and ♀ flowers often, in catkins. Perianth absent or haplo- chlamydeous and bracteoid</i>		
Order 1. Salicales	33	157
Fam. SALICACEAE	33	158
Order 2. Myricales	34	159
Fam. MYRICACEAE	34	159
Order 3. *Juglandales	34	160
Fam. *JUGLANDACEAE	34	160
Order 4. Fagales	34	161
Fam. 1. BETULACEAE	35	162
Fam. 2. FAGACEAE	35	162
Order 5. Urticales	36	164
Fam. 1. ULMACEAE	36	164
Fam. 2. CANNABACEAE (<i>Moraceae, Can- naboidae in Syllabus</i>)	36	167
Fam. 3. URTICACEAE	37	167

Synoptic Index

xiii

<i>b. Petaloideae (Moss). Flowers haplo-</i>		
<i>chlamydeous, perianth often</i>		
<i>petaloid. Diplochlamydeous</i>		
	<i>flowers rare</i>	37 168
Order 6	Santalales	37 170
	Fam. 1. SANTALACEAE	38 170
	Fam. 2. LORANTHACEAE	38 172
Order 7.	Aristolochiales	38 176
	Fam. ARISTOLOCHIACEAE	38 176
Order 8.	Polygonales	39 177
	Fam. POLYGONACEAE	39 177
<i>c. (Consisting of the one order Centro-</i>		
<i>spermae.) Corolla usually</i>		
<i>haplochlamydeous, sepaloid</i>		
<i>or petaloid. Heterochlamydy,</i>		
<i>however, is not uncommon</i>		
Order 9.	Centrospermae	40 179
	Fam. 1. CHENOPODIACEAE	40 179
	Fam. 2. *AMARANTACEAE	41 181
	Fam. 3. *AIZOACEAE	42 183
	Fam. 4. PORTULACACEAE	42 183
	Fam. 5. CARYOPHYLLACEAE	43 184
<i>d. Heterochlamydeae (Moss). Orders with</i>		
<i>predominant heterochlamydy</i>		
	<i>a. Apocarpy and Hypogyny predominant</i>	45 185
Order 10.	Ranales	45 185
	Fam. 1. NYMPHAEACEAE	45 185
	Fam. 2. CERATOPHYLLACEAE	46 186
	Fam. 3. RANUNCULACEAE	46 187
	Fam. 4. BERBERIDACEAE	48 188
	<i>β. Syncarpy and Hypogyny predominant</i>	48 195
Order 11.	Rhoeadales	49 195
	Fam. 1. PAPAVERACEAE	49 195
	Fam. 2. CRUCIFERAE	50 197
	Fam. 3. RESEDACEAE	54 201

Order 12.	Sarraceniales	.	.	.	.	54	201
	Fam. DROSERACEAE	.	.	.	.	54	202
7.	<i>Apocarpy and Hypogyny still occur but Perigyny becomes more frequent. By sinking of the Gynaecium into the hollow flower-axis Syncarpy and Epigynous insertion of the perianth and stamens also takes place</i>					54	203
Order 13.	Rosales	.	.	.	.	54	203
	Fam. 1. CRASSULACEAE	.	.	.	.	55	204
	Fam. 2. SAXIFRAGACEAE	.	.	.	.	55	204
	Fam. 3. *PLATANACEAE	.	.	.	.	56	209
	Fam. 4. ROSACEAE	.	.	.	.	57	210
	Fam. 5. LEGUMINOSAE	.	.	.	.	59	215
8.	<i>The flowers have predominantly 5 or 4 whorls. Apocarpy and Isomery still occur, but Syncarpy and Oligomery of the Gynaecium preponderate. Pleiomery of Gynaecium rare</i>					61	223
Order 14.	Geraniales	.	.	.	.	61	223
	Fam. 1. GERANIACEAE	.	.	.	.	62	223
	Fam. 2. OXALIDACEAE	.	.	.	.	63	224
	Fam. 3. LINACEAE	.	.	.	.	63	226
	Fam. 4. POLYGALACEAE	.	.	.	.	63	235
	Fam. 5. EUPHORBIACEAE	.	.	.	.	63	236
	Fam. 6. CALLITRICHACEAE	.	.	.	.	64	241
Order 15.	Sapindales	.	.	.	.	64	241
	Fam. 1. BUXACEAE	.	.	.	.	65	241
	Fam. 2. EMPETRACEAE	.	.	.	.	65	241
	Fam. 3. AQUIFOLIACEAE	.	.	.	.	65	244
	Fam. 4. CELASTRACEAE	.	.	.	.	65	245
	Fam. 5. *STAPHYLEACEAE	.	.	.	.	66	246
	Fam. 6. ACERACEAE	.	.	.	.	66	246
	Fam. 7. *HIPPOCASTANACEAE	.	.	.	.	66	247
	Fam. 8. BALSAMINACEAE	.	.	.	.	67	249

Synoptic Index

xv

Order 16. Rhamnales	67	250
Fam. RHAMNACEAE	67	250
Order 17. Malvales	67	252
Fam. 1. TILIACEAE	68	253
Fam. 2. MALVACEAE	68	253
e. <i>Flowers spirocyclic or of 5-4 whorls. Syncarpy the rule (Apocarpy in the more primitive, exotic families). Tendency for gynaeceum to become sunk in flower-axis</i>		
	69	257
Order 18. Parietales	69	257
Fam. 1. GUTTIFERAE	69	261
Fam. 2. ELATINACEAE	70	264
Fam. 3. FRANKENIACEAE	70	264
Fam. 4. *TAMARICACEAE	70	264
Fam. 5. CISTACEAE	71	266
Fam. 6. VIOLACEAE	71	266
f. <i>The flowers are cyclic and the sinking of the gynaeceum into the hollow flower-axis is general: connation of gynaeceum and flower-axis predominates</i>		
	71	274
Order 19. Myrtiflorae	71	274
Fam. 1. THYMELAEACEAE	72	275
Fam. 2. ELAEAGNACEAE	72	276
Fam. 3. LYTHRACEAE	72	276
Fam. 4. EPILOBIACEAE (Oenotheraceae)	73	283
Fam. 5. HALORRHAGACEAE	74	284
Fam. 6. HIPPURIDACEAE	74	285
Order 20. Umbellales	74	285
Fam. 1. ARALIACEAE	74	285
Fam. 2. UMBELLIFERAE	75	286
Fam. 3. CORNACEAE	80	290
Subclass 2. METACHLAMYPDEAE	80	291
A. Polypetal, as well as sympetal, occurs. Two whorls or one		

	whorl of stamens. Hypogyny predominates, but epigyny also occurs	80	291
Order 1.	Ericales	80	291
	Fam. 1. PIROLACEAE	80	291
	Fam. 2. ERICACEAE	81	292
Order 2.	Primulales	83	295
	Fam. PRIMULACEAE	83	295
Order 3.	Plumbaginales	84	296
	Fam. PLUMBAGINACEAE	85	296
B. (C. in Syllabus).	Sympetaly dominant. Stamens always in one whorl. Union of Carpels sometimes imperfect. Hypogyny usual.	85	300
Order 4.	Contortae	85	300
	Fam. 1. OLEACEAE	85	300
	Fam. 2. GENTIANACEAE	86	303
	Fam. 3. APOCYNACEAE	87	303
C. (D. in Syllabus).	Sympetaly constant. Only one whorl of stamens present. Carpels usually 2, always perfectly united. Zygomorphy frequent	87	308
a.	<i>Perianth hypogynous</i>	87	308
Order 5.	Tubiflorae	88	308
	Fam. 1. CONVULVACEAE	88	308
	Fam. 2. POLEMONIACEAE	88	310
	Fam. 3. BORRAGINACEAE	89	311
	Fam. 4. VERBENACEAE	91	313
	Fam. 5. LABIATAE	91	315
	Fam. 6. SOLANACEAE	94	320
	Fam. 7. SCROPHULARIACEAE	95	322
	Fam. 8. OROBANCHACEAE	97	327
	Fam. 9. LENTIBULARIACEAE	97	329

Synoptic Index

xvii

Order 6. Plantaginales		98	332
Fam. PLANTAGINACEAE		98	332
<i>b. Perianth epigynous</i>		98	333
<i>a. Stamens free</i>		98	333
Order 7. Rubiales		99	333
Fam. 1. RUBIACEAE		99	333
Fam. 2. CAPRIFOLIACEAE		99	336
Fam. 3. ADOXACEAE		100	337
Fam. 4. VALERIANACEAE		101	337
Fam. 5. DIPSACACEAE		101	338
<i>β. Stamens conniving or partly united</i>		101	338
Order 8. Cucurbitales		102	338
Fam. CUCURBITACEAE		102	338
Order 9. Campanulatae		102	340
Fam. 1. CAMPANULACEAE		102	340
Fam. 2. COMPOSITAE		103	342