

Contents

<i>Advisory board</i>	xiv
<i>About the authors</i>	xv
<i>Acknowledgements</i>	xvi
1. About this book	1
2. Habitat protection	7
Key messages	7
2.1. Legally protect habitats	8
2.2. Ensure connectivity between habitat patches	9
2.3. Provide or retain un-harvested buffer strips	10
3. Education and awareness raising	13
Key messages	13
3.1. Raise awareness amongst the general public through campaigns and public information	13
3.2. Provide bird feeding materials to families with young children	14
3.3. Enhance bird taxonomy skills through higher education and training	14
3.4. Provide training to conservationists and land managers on bird ecology and conservation	14
4. Threat: Residential and commercial development	15
Key messages	15
4.1. Angle windows to reduce collisions	15
4.2. Mark or tint windows to reduce collision mortality	15
5. Threat: Agriculture	17
Key messages – All farming systems	17
Key messages – Arable farming	20
Key messages – Livestock farming	21
Key messages – Perennial, non-timber crops	23
Key messages – Aquaculture	24
All farming systems	25
5.1. Support or maintain low-intensity agricultural systems	25
5.2. Practise integrated farm management	25
5.3. Food labelling schemes relating to biodiversity-friendly farming	25
5.4. Increase the proportion of natural/semi-natural vegetation in the farmed landscape	25
5.5. Pay farmers to cover the costs of conservation measures	27
5.6. Cross compliance standards for all subsidy payments	36
5.7. Reduce field size (or maintain small fields)	37
5.8. Provide or retain set-aside areas in farmland	37
5.9. Manage hedges to benefit wildlife	42
5.10. Plant new hedges	44
5.11. Manage stone-faced hedge banks to benefit birds	44
5.12. Manage ditches to benefit wildlife	45
5.13. Protect in-field trees	46
5.14. Plant in-field trees	46

5.15.	Tree pollarding and tree surgery	46
5.16.	Plant wild bird seed or cover mixture	46
5.17.	Plant nectar flower mixture/wildflower strips	55
5.18.	Create uncultivated margins around intensive arable or pasture fields	57
5.19.	Plant grass buffer strips/margins around arable or pasture fields	59
5.20.	Use mowing techniques to reduce chick mortality	63
5.21.	Provide refuges in fields during harvest or mowing	64
5.22.	Mark bird nests during harvest or mowing	64
5.23.	Relocate nests at harvest time to reduce nestling mortality	65
5.24.	Make direct payments per clutch for farmland birds	65
5.25.	Control scrub on farmland	66
5.26.	Take field corners out of management	67
5.27.	Reduce conflict by deterring birds from taking crops	67
Arable farming		68
5.28.	Increase crop diversity	68
5.29.	Implement 'mosaic management'	69
5.30.	Leave overwinter stubbles	69
5.31.	Plant nettle strips	73
5.32.	Leave unharvested cereal headlands within arable fields	73
5.33.	Plant crops in spring rather than autumn	73
5.34.	Undersow spring cereals, with clover for example	75
5.35.	Plant more than one crop per field (intercropping)	76
5.36.	Revert arable land to permanent grassland	76
5.37.	Reduce tillage	78
5.38.	Add 1% barley into wheat crop for corn buntings	81
5.39.	Leave uncropped, cultivated margins or plots (includes lapwing and stone curlew plots)	81
5.40.	Create skylark plots	83
5.41.	Create corn bunting plots	84
5.42.	Plant cereals in wide-spaced rows	85
5.43.	Create beetle banks	85
Livestock farming		87
5.44.	Maintain species-rich, semi-natural grassland	87
5.45.	Reduce management intensity on permanent grasslands	88
5.46.	Reduce grazing intensity	90
5.47.	Provide short grass for waders	93
5.48.	Raise mowing height on grasslands	93
5.49.	Delay mowing date or first grazing date on grasslands	94
5.50.	Leave uncut rye grass in silage fields	95
5.51.	Plant cereals for wholecrop silage	96
5.52.	Maintain lowland heathland	97
5.53.	Maintain rush pastures	97
5.54.	Maintain traditional water meadows	98
5.55.	Maintain upland heath/moor	99
5.56.	Plant Brassica fodder crops	100
5.57.	Use mixed stocking	100
5.58.	Use traditional breeds of livestock	100
5.59.	Maintain wood pasture and parkland	100
5.60.	Exclude grazers from semi-natural habitats (including woodland)	100
5.61.	Protect nests from livestock to reduce trampling	103
5.62.	Mark fences to reduce bird collision mortality	103
5.63.	Create open patches or strips in permanent grassland	104

Perennial, non-timber crops	104
5.64. Maintain traditional orchards	105
5.65. Manage perennial bioenergy crops to benefit wildlife	105
Aquaculture	106
5.66. Reduce conflict with humans to reduce persecution	106
5.67. Scare birds from fish farms	106
5.68. Disturb birds at roosts	109
5.69. Use electric fencing to exclude fish-eating birds	110
5.70. Use netting to exclude fish-eating birds	111
5.71. Disturb birds using foot patrols	112
5.72. Use 'mussel socks' to prevent birds from attacking shellfish	112
5.73. Translocate birds away from fish farms	113
5.74. Increase water turbidity to reduce fish predation by birds	113
5.75. Provide refuges for fish within ponds	114
5.76. Use in-water devices to reduce fish loss from ponds	114
5.77. Spray water to deter birds from ponds	115
5.78. Deter birds from landing on shellfish culture gear	115
6. Threat: Energy production and mining	117
Key messages	117
6.1. Paint wind turbines to increase their visibility	117
7. Threat: Transportation and service corridors	118
Key messages – Verges and airports	118
Key messages – Power lines and electricity pylons	118
Verges and airports	119
7.1. Mow roadside verges	119
7.2. Sow roadside verges	119
7.3. Scare or otherwise deter birds from airports	119
Power lines and electricity pylons	120
7.4. Bury or isolate power lines to reduce incidental mortality	120
7.5. Remove earth wires to reduce incidental mortality	121
7.6. Thicken earth wire to reduce incidental mortality	121
7.7. Mark power lines to reduce incidental mortality	121
7.8. Use raptor models to deter birds and so reduce incidental mortality	124
7.9. Add perches to electricity pylons to reduce electrocution	124
7.10. Insulate power pylons to prevent electrocution	125
7.11. Use perch-deterrents to stop raptors perching on pylons	125
7.12. Reduce electrocutions by using plastic, not aluminium, leg rings to mark birds	126
8. Threat: Biological resource use	127
Key messages – reducing exploitation and conflict	127
Key messages – reducing fisheries bycatch	128
Reducing exploitation and conflict	130
8.1. Use legislative regulation to protect wild populations	130
8.2. Increase 'on-the-ground' protection to reduce unsustainable levels of exploitation	132
8.3. Promote sustainable alternative livelihoods	133
8.4. Use education programmes and local engagement to help reduce persecution or exploitation of species	133
8.5. Employ local people as 'biomonitors'	135

8.6.	Mark eggs to reduce their appeal to egg collectors	136
8.7.	Relocate nestlings to reduce poaching	136
8.8.	Use wildlife refuges to reduce hunting disturbance	137
8.9.	Introduce voluntary 'maximum shoot distances'	138
8.10.	Provide 'sacrificial' grasslands to reduce the impact of wild geese on crops	138
8.11.	Move fish-eating birds to reduce conflict with fishermen	139
8.12.	Scare fish-eating birds from areas to reduce conflict	139
	Reduce fisheries bycatch	139
8.13.	Set longlines at night to reduce seabird bycatch	140
8.14.	Turn deck lights off during night-time setting of longlines to reduce bycatch	142
8.15.	Use streamer lines to reduce seabird bycatch on longlines	143
8.16.	Use larger hooks to reduce seabird bycatch	146
8.17.	Use a water cannon when setting longlines to reduce seabird bycatch	147
8.18.	Set lines underwater to reduce seabird bycatch	147
8.19.	Set longlines at the side of the boat to reduce seabird bycatch	148
8.20.	Use a line shooter to reduce seabird bycatch	148
8.21.	Use bait throwers to reduce seabird bycatch	149
8.22.	Tow buoys behind longlining boats to reduce seabird bycatch	149
8.23.	Dye baits to reduce seabird bycatch	149
8.24.	Use high-visibility longlines to reduce seabird bycatch	150
8.25.	Use a sonic scarer when setting longlines to reduce seabird bycatch	150
8.26.	Weight baits or lines to reduce longline bycatch of seabirds	150
8.27.	Use shark liver oil to deter birds when setting lines	152
8.28.	Thaw bait before setting lines to reduce seabird bycatch	152
8.29.	Reduce seabird bycatch by releasing offal overboard when setting longlines	153
8.30.	Use bird exclusion devices such as 'Brickle curtains' to reduce seabird mortality when hauling longlines	153
8.31.	Use acoustic alerts on gillnets to reduce seabird bycatch	154
8.32.	Use high-visibility mesh on gillnets to reduce seabird bycatch	154
8.33.	Reduce gillnet deployment time to reduce seabird bycatch	155
8.34.	Mark trawler warp cables to reduce seabird collisions	155
8.35.	Reduce 'ghost fishing' by lost/discarded gear	155
8.36.	Reduce bycatch through seasonal or area closures	156
9.	Threat: Human intrusions and disturbance	157
	Key messages	157
9.1.	Use wildlife refuges to reduce hunting disturbance	157
9.2.	Use signs and access restrictions to reduce disturbance at nest sites	157
9.3.	Set minimum distances for approaching birds (buffer zones)	160
9.4.	Provide paths to limit the extent of disturbance	160
9.5.	Reduce visitor group sizes	161
9.6.	Use voluntary agreements with local people to reduce disturbance	161
9.7.	Start educational programmes for personal watercraft owners	161
9.8.	Habituate birds to human visitors	162
9.9.	Use nest covers to reduce the impact of research on predation of ground-nesting seabirds	162
10.	Threat: Natural system modifications	164
	Key messages	164
10.1.	Use prescribed burning	167
10.2.	Use fire suppression/control	183

10.3.	Protect nest trees before burning	184
10.4.	Clear or open patches in forests	184
10.5.	Clearcut and re-seed forests	186
10.6.	Thin trees within forests	187
10.7.	Coppice trees	189
10.8.	Use patch retention harvesting instead of clearcutting	190
10.9.	Use selective harvesting/logging instead of clearcutting	191
10.10.	Use variable retention management during forestry operations	192
10.11.	Use shelterwood cutting instead of clearcutting	193
10.12.	Manage woodland edges for birds	193
10.13.	Manually control or remove mid-storey and ground-level vegetation (including mowing, chaining, cutting etc)	194
10.14.	Replace non-native species of tree/shrub	202
10.15.	Provide deadwood/snags in forests	202
10.16.	Remove coarse woody debris from forests	204
10.17.	Apply herbicide to mid- and under-storey vegetation	205
10.18.	Treat wetlands with herbicide	206
10.19.	Employ grazing in natural and semi-natural habitats	208
10.20.	Plant trees to act as windbreaks	214
10.21.	Re-seed grasslands	215
10.22.	Fertilise artificial grasslands	215
10.23.	Raise water levels in ditches or grassland	217
10.24.	Manage water level in wetlands	218
10.25.	Use environmentally sensitive flood management	220
10.26.	Use greentree reservoir management	220
10.27.	Plough habitats	221
10.28.	Create scrapes and pools in wetlands and wet grasslands	222
11.	Habitat restoration and creation	224
	Key messages	224
11.1.	Restore or create forests	225
11.2.	Restore or create grasslands	229
11.3.	Restore or create traditional water meadows	235
11.4.	Restore or create shrubland	236
11.5.	Restore or create savannas	237
11.6.	Restore or create wetlands and marine habitats	237
12.	Threat: Invasive alien and other problematic species	243
	Key messages – reduce predation by other species	243
	Key messages – reduce incidental mortality during predator eradication or control	244
	Key messages – reduce nest predation by excluding predators from nests or nesting areas	244
	Key messages – reduce mortality by reducing hunting ability or changing predator behaviour	245
	Key messages – reduce competition with other species for food and nest sites	246
	Key messages – reduce adverse habitat alteration by other species	246
	Key messages – reduce parasitism and disease	247
	Key messages – reduce detrimental impacts of other problematic species	247
	Reduce predation by other species	248
12.1.	Remove or control predators to enhance bird populations and communities	248

Predator control on islands	249
12.2. Control avian predators on islands	249
12.3. Control mammalian predators on islands	251
12.4. Control invasive ants on islands	261
12.5. Control predators not on islands	261
12.6. Reduce predation by translocating predators	269
Reduce incidental mortality during predator eradication or control	270
12.7. Do birds take bait designed for pest control?	270
12.8. Distribute poison bait using dispensers	271
12.9. Use repellents on baits	271
12.10. Use coloured baits to reduce accidental mortality during predator control	272
Reduce nest predation by excluding predators from nests or nesting areas	273
12.11. Physically protect nests from predators using non-electric fencing	273
12.12. Protect bird nests using electric fencing	274
12.13. Physically protect nests with individual exclosures/barriers or provide shelters for chicks	276
12.14. Can nest protection increase nest abandonment?	281
12.15. Can nest protection increase predation of adults and chicks?	282
12.16. Use artificial nests that discourage predation	283
12.17. Use multiple barriers to protect nests	284
12.18. Plant nesting cover to reduce nest predation	285
12.19. Use snakeskin to deter mammalian nest predators	285
12.20. Use mirrors to deter nest predators	285
12.21. Use naphthalene to deter mammalian predators	286
12.22. Use ultrasonic devices to deter cats	286
12.23. Protect nests from ants	286
12.24. Guard nests to prevent predation	287
12.25. Use 'cat curfews' to reduce predation	287
12.26. Use lion dung to deter domestic cats	287
12.27. Play spoken-word radio programmes to deter predators	287
Reduce mortality by reducing hunting ability or changing predator behaviour	288
12.28. Use collar-mounted devices to reduce predation	288
12.29. Use supplementary feeding to reduce predation	288
12.30. Use aversive conditioning to reduce nest predation	289
12.31. Reduce predation by translocating nest boxes	293
Reduce competition with other species for food and nest sites	294
12.32. Reduce inter-specific competition for nest sites by removing competitor species	294
12.33. Reduce inter-specific competition for nest sites by modifying habitats to exclude competitor species	298
12.34. Protect nest sites from competitors	298
12.35. Reduce competition between species by providing nest boxes	299
12.36. Reduce inter-specific competition for food by removing or controlling competitor species	300
Reduce adverse habitat alteration by other species	301
12.37. Reduce adverse habitat alterations by excluding problematic species	301
12.38. Control or remove habitat-altering mammals	303
12.39. Remove problematic vegetation	304
12.40. Use buffer zones to reduce the impact of invasive plant control	305

Reduce parasitism and disease	306
12.41. Remove/treat endoparasites and diseases	306
12.42. Exclude or control 'reservoir species' to reduce parasite burdens	307
12.43. Remove/treat ectoparasites to increase survival or reproductive success	308
12.44. Guard nests to reduce risk of ectoparasites	312
12.45. Remove/control brood parasites	312
12.46. Use false brood parasite eggs to discourage brood parasitism	316
12.47. Provide supplementary food to increase parental presence and so reduce brood parasitism	317
12.48. Alter artificial nest sites to discourage brood parasitism	317
Reduce detrimental impacts of other problematic species	317
12.49. Use copper strips to exclude snails from nests	317
13. Threat: Pollution	318
Key messages – Industrial pollution	318
Key messages – Agricultural pollution	318
Key message – Air-borne pollutants	319
Key messages – Excess energy	319
Industrial pollution	320
13.1. Clean birds following oil spills	320
13.2. Relocate birds following oil spills	322
13.3. Deter or prevent birds from landing on toxic pools	323
13.4. Use repellents to deter birds from landing on pools polluted by mining	325
Agricultural pollution	325
13.5. Reduce pesticide or herbicide use generally	325
13.6. Restrict certain pesticides or other agricultural chemicals	327
13.7. Provide food for vultures to reduce mortality from diclofenac	328
13.8. Make selective use of spring herbicides	329
13.9. Use organic rather than mineral fertilisers	329
13.10. Reduce chemical inputs in permanent grassland management	329
13.11. Leave headlands in fields unsprayed (conservation headlands)	330
13.12. Provide unfertilised cereal headlands in arable fields	332
13.13. Plant riparian buffer strips	332
13.14. Provide buffer strips around in-field ponds	333
Air-borne pollutants	333
13.15. Use lime to reduce acidification in lakes	333
Excess energy	333
13.16. Reduce incidental mortality from birds being attracted to artificial lights	333
13.17. Turn off lights to reduce mortality from artificial lights	334
13.18. Reduce the intensity of lighthouse beams	334
13.19. Shield lights to reduce mortality from artificial lights	334
13.20. Use flashing lights to reduce mortality from artificial lights	335
13.21. Use lights low in spectral red to reduce mortality from artificial lights	335
13.22. Use volunteers to collect downed birds and rehabilitate them	336
14. Threat: Climate change, extreme weather and geological events	337
Key messages	337
14.1. Water nesting mounds to increase incubation success in malleefowl	337
14.2. Replace nesting substrate following severe weather	338

15. General responses to small/declining populations	339
Key messages – inducing breeding, rehabilitation and egg removal	339
Key messages – provide artificial nesting sites	339
Key messages – foster chicks in the wild	340
Key messages – provide supplementary food	341
Key messages – translocations	342
15.1. Use artificial visual and auditory stimuli to induce breeding in wild populations	343
15.2. Rehabilitate injured birds	343
15.3. Remove eggs from wild nests to increase reproductive output	344
15.4. Provide artificial nesting sites	345
15.5. Clean nest boxes to increase occupancy or reproductive success	390
15.6. Use differently-coloured artificial nests	392
15.7. Provide nesting material for wild birds	393
15.8. Repair/support nests to support breeding	393
15.9. Artificially incubate eggs or warm nests	394
15.10. Provide nesting habitat for birds that is safe from extreme weather	394
15.11. Remove vegetation to create nesting areas	395
15.12. Guard nests to increase nest success	397
Foster chicks in the wild	398
15.13. Foster eggs or chicks with wild conspecifics	398
15.14. Foster eggs or chicks with wild non-conspecifics (cross-fostering)	404
Provide supplementary food	407
15.15. Provide supplementary food to increase reproductive success	408
15.16. Provide supplementary food to allow the rescue of a second chick	430
15.17. Provide supplementary food to increase adult survival	431
15.18. Can supplementary feeding increase predation or parasitism?	449
15.19. Provide supplementary food through the establishment of food populations	450
15.20. Use perches to increase foraging success	451
15.21. Place feeders close to windows to reduce collisions	452
15.22. Provide supplementary water to increase survival or reproductive success	453
15.23. Provide calcium supplements to increase survival or reproductive success	453
Translocations	457
15.24. Translocate birds to re-establish populations or increase genetic variation	457
15.25. Use techniques to increase the survival of species after capture	474
15.26. Ensure translocated birds are familiar with each other before release	475
15.27. Ensure genetic variation to increase translocation success	475
15.28. Translocate nests to avoid disturbance	476
15.29. Use vocalisations to attract birds to new sites	477
15.30. Use decoys to attract birds to new sites	479
15.31. Alter habitat to encourage birds to leave an area	481
16. Captive breeding, rearing and releases (<i>ex situ</i> conservation)	483
Key messages – captive breeding	483
Key messages – release of captive-bred individuals	484

Captive breeding	485
16.1. Use captive breeding to increase or maintain populations	485
16.2. Can captive breeding have deleterious effects on individual fitness?	490
16.3. Use artificial insemination in captive breeding	491
16.4. Freeze semen for use in artificial insemination	492
16.5. Wash contaminated semen and use it for artificial insemination	493
16.6. Artificially incubate and hand-rear birds in captivity	494
16.7. Use puppets to increase the success of hand-rearing	504
Release of captive-bred individuals	505
16.8. Release captive-bred individuals into the wild to restore or augment wild populations	505
16.9. Use appropriate populations to source released populations	520
16.10. Use holding pens at release sites	521
16.11. Clip birds' wings on release	523
16.12. Release birds in groups	524
16.13. Release chicks and adults in 'coveys'	524
16.14. Release birds as adults or sub-adults, not juveniles	525
16.15. Use 'anti-predator training' to improve survival after release	527
16.16. Use 'flying training' before release	528
16.17. Provide supplementary food during and after release	529
16.18. Use microlites to help birds migrate	530
<i>Index</i>	531