

brief contents

Introduction 1

1 A leaf cell consists of several metabolic compartments 3

- 1.1 The cell wall gives the plant cell mechanical stability 5
 - 1.2 Vacuoles have multiple functions 10
 - 1.3 Plastids have evolved from cyanobacteria 12
 - 1.4 Mitochondria also result from endosymbionts 17
 - 1.5 Peroxisomes are the site of reactions in which toxic intermediates are formed 18
 - 1.6 The endoplasmic reticulum and Golgi apparatus form a network for the distribution of biosynthesis products 19
 - 1.7 Functionally intact cell organelles can be isolated from plant cells 21
 - 1.8 Various transport processes facilitate the exchange of metabolites between different compartments 24
 - 1.9 Translocators catalyse the specific transport of substrates and products of metabolism 25
 - 1.10 Ion channels have a very high transport capacity 31
 - 1.11 Porins consist of β -sheet structures 34
- Further reading 37

2 The use of energy from sunlight by photosynthesis 39

- 2.1 How did photosynthesis start? 39
 - 2.2 Pigments capture energy from sunlight 41
 - 2.3 Light absorption excites the chlorophyll molecule 46
 - 2.4 An antenna is required to capture light 50
- Further reading 59

3 Photosynthesis is an electron transport process 60

- 3.1 The photosynthetic machinery is constructed from modules 61
- 3.2 A reductant and an oxidant are formed during photosynthesis 64
- 3.3 The basic structure of a photosynthetic reaction centre has been resolved by X-ray structure analysis 65
- 3.4 How does a reaction centre function? 69
- 3.5 Two photosynthetic reaction centres are arranged in tandem in photosynthesis of algae and plants 73

- 3.6 Water is split by photosystem II 76
- 3.7 The cytochrome- b_6/f complex mediates electron transport between photosystem II and photosystem I 84
- 3.8 Photosystem I reduces NADP 91
- 3.9 In the absence of other acceptors electrons can be transferred from PS I to oxygen 96
- 3.10 Regulatory processes control the distribution of the captured excitons between the two photosystems 99
- Further reading 103
- 4 ATP generation by photosynthesis 105**
 - 4.1 A proton gradient serves as an energy-rich intermediate state during ATP synthesis 106
 - 4.2 The electrochemical proton gradient can be dissipated by uncouplers 108
 - 4.3 H^+ -ATP synthases from bacteria, chloroplasts, and mitochondria have a common basic structure 111
 - 4.4 The synthesis of ATP is effected by a conformational change of the protein 116
 - Further reading 120
- 5 Mitochondria, the power stations of the cell 121**
 - 5.1 Biological oxidation is preceded by substrate degradation 122
 - 5.2 Mitochondria are the sites of cell respiration 122
 - 5.3 Degradation of substrates for biological oxidation takes place in the matrix compartment 124
 - 5.4 How much energy can be gained by the oxidation of NADH? 131
 - 5.5 The mitochondrial respiratory chain shares common features with the photosynthetic electron transport chain 132
 - 5.6 Electron transport of the respiratory chain is coupled via proton transport to the synthesis of ATP 138
 - 5.7 Plant mitochondria have special metabolic functions 142
 - 5.8 Compartmentalization of mitochondrial metabolism requires specific membrane translocators 145
 - Further reading 146
- 6 Photosynthetic CO_2 assimilation by the Calvin cycle 148**
 - 6.1 CO_2 assimilation proceeds via the dark reaction of photosynthesis 148
 - 6.2 Ribulose biphosphate carboxylase catalyses the fixation of CO_2 151
 - 6.3 The reduction of 3-phosphoglycerate yields triose phosphate 156
 - 6.4 Ribulose biphosphate is regenerated from triose phosphate 158
 - 6.5 Besides the reductive pentose phosphate pathway there is also an oxidative pentose phosphate pathway 161
 - 6.6 Reductive and oxidative pentose phosphate pathways are regulated 167
 - Further reading 174

7 Photorespiration 175

- 7.1 Ribulose 1,5-bisphosphate is recovered by recycling 2-phosphoglycolate 177
 - 7.2 The NH_4^+ released in the photorespiratory pathway is refixed in the chloroplasts 182
 - 7.3 Peroxisomes have to be provided with reducing equivalents to reduce hydroxypyruvate 183
 - 7.4 The peroxisomal matrix is a special compartment for the interconversion of toxic products 187
 - 7.5 How high are the costs of the ribulose bisphosphate oxygenase reaction for the plant? 188
 - 7.6 There is no net CO_2 fixation at the compensation point 189
 - 7.7 The photorespiration pathway, although energy-consuming, may also have a useful function for the plant 190
- Further reading 190

8 Photosynthesis and water consumption 192

- 8.1 The entry of CO_2 into the leaf is accompanied by an escape of water vapour from the leaves 192
 - 8.2 Stomata regulate the gas exchange of a leaf 194
 - 8.3 The diffusive flux of CO_2 into a plant cell 198
 - 8.4 C_4 plants perform CO_2 assimilation with less water consumption 200
 - 8.5 Crassulacean acid metabolism makes it possible for plants to survive even during very severe water shortage 213
- Further reading 218

9 Polysaccharides 219

- 9.1 Large quantities of carbohydrate can be stored as starch in the cell 220
 - 9.2 Sucrose synthesis takes place in the cytosol 229
 - 9.3 Utilization of triose phosphate formed during photosynthesis is strictly regulated 231
 - 9.4 In some plants assimilates from the leaves are exported as sugar alcohols or oligosaccharides of the raffinose family 238
 - 9.5 Fructans are deposited as storage substances in the vacuole 239
 - 9.6 Cellulose is synthesized by enzymes located in the plasma membrane 243
- Further reading 246

10 Nitrate assimilation 247

- 10.1 The reduction of nitrate to NH_3 proceeds in two partial reactions 248
- 10.2 Nitrate assimilation also takes place in the roots 254

- 10.3 Nitrate assimilation is strictly controlled 255
- 10.4 The end-product of nitrate assimilation is a whole spectrum of amino acids 258
- 10.5 Glutamate is the precursor for synthesis of chlorophylls and cytochromes 271
- Further reading 276

11 Nitrogen fixation 277

- 11.1 Legumes form a symbiosis with nodule-forming bacteria 278
- 11.2 N_2 fixation can proceed only at very low oxygen concentrations 286
- 11.3 The energy costs for the utilization of N_2 as a nitrogen source are much higher than for the utilization of NO_3^- 288
- Further reading 288

12 Sulfate assimilation 289

- 12.1 Sulfate assimilation proceeds by photosynthesis 289
- 12.2 Glutathione serves the cell as an antioxidant and is an agent for the detoxification of pollutants 294
- 12.3 Methionine is synthesized from cysteine 297
- 12.4 Excessive concentrations of sulfur dioxide in air are toxic for plants 299
- Further reading 300

13 Phloem transport 301

- 13.1 There are two paths for phloem loading 303
- 13.2 Phloem transport proceeds by mass flow 305
- 13.3 Sink tissues are supplied by phloem unloading 306
- Further reading 308

14 Plant storage proteins 309

- 14.1 Globulins are the most abundant storage proteins 310
- 14.2 Prolamins are formed as storage proteins in grasses 311
- 14.3 2S-Proteins are present in seeds of dicotyledonous plants 311
- 14.4 Special proteins protect seeds from being eaten by animals 312
- 14.5 Synthesis of the storage proteins occurs on the rough endoplasmic reticulum 313
- 14.6 Proteinases mobilize the amino acids deposited in storage proteins 315
- Further reading 315

15 Glycerolipids 317

- 15.1 Polar glycerolipids are membrane constituents 317
- 15.2 Triacylglycerols are storage substances 323
- 15.3 The *de novo* synthesis of fatty acids takes place in the plastids 324
- 15.4 Glycerol 3-phosphate is the precursor for the synthesis of glycerolipids 334
- 15.5 Triacylglycerols are formed in the membranes of the endoplasmic reticulum 338

- 15.6 During seed germination storage lipids are mobilized for the production of carbohydrates in the glyoxysomes 343
- 15.7 Lipoxygenase is involved in the synthesis of aromatic defence and signal substances 347
- Further reading 350

16 The function of secondary metabolites in plants 352

- 16.1 Secondary metabolites often protect plants from pathogenic micro-organisms and herbivores 352
- 16.2 Alkaloids comprise a variety of heterocyclic secondary metabolites 354
- 16.3 Some plants emit prussic acid when wounded by animals 356
- 16.4 When wounded, some plants emit volatile mustard oils 357
- 16.5 Plants protect themselves by tricking herbivores with false amino acids 357
- Further reading 358

17 Isoprenoids 360

- 17.1 Acetyl CoA is the precursor for the synthesis of isoprenoids 362
- 17.2 Prenyl transferases catalyse the association of isoprene units 364
- 17.3 Some plants emit isoprenes into the air 365
- 17.4 Many aromatic substances are derived from geranyl pyrophosphate 367
- 17.5 Farnesyl pyrophosphate is the precursor for the formation of sesquiterpenes 368
- 17.6 Geranylgeranyl pyrophosphate is the precursor for defence substances, phytohormones, and carotenoids 371
- 17.7 A prenyl chain renders substances lipid soluble 372
- 17.8 The regulation of isoprenoid synthesis 374
- 17.9 Isoprenoids are very stable and persistent substances 376
- Further reading 376

18 Phenylpropanoids 377

- 18.1 Phenylalanine ammonia lyase catalyses the initial reaction of phenylpropanoid metabolism 379
- 18.2 Mono-oxygenases are involved in the synthesis of phenols 379
- 18.3 Phenylpropanoid compounds polymerize to macromolecules 382
- 18.4 For the synthesis of flavonoids and stilbenes a second aromatic ring is formed from acetate residues 387
- 18.5 Flavonoids have multiple functions in plants 388
- 18.6 Anthocyanins are flower pigments and protect plants against excessive light 389
- 18.7 Tannins bind tightly to proteins and therefore have defence functions 391
- Further reading 393

19 Signals regulating the growth and development of plant organs 394

- 19.1 Auxin stimulates shoot elongation 396

- 19.2 Gibberellins regulate stem elongation 398
- 19.3 Cytokinins stimulate cell division 400
- 19.4 Absciscic acid controls the water balance of the plant 401
- 19.5 Ethylene makes fruit ripen 402
- 19.6 Phytochromes function as light sensors 403
- 19.7 Signal transduction chains known from animal metabolism could be models for signal transduction in plants 406
- Further reading 413

20 The genomes of plant cells 415

- 20.1 In the nucleus the genetic information is divided among several chromosomes 416
- 20.2 The DNA of the nuclear genome is transcribed by three specialized RNA polymerases 419
- 20.3 DNA polymorphism yields genetic markers for plant breeding 428
- 20.4 Transposable DNA elements roam through the genome 433
- 20.5 Most plant cells contain viruses 435
- 20.6 Plastids possess a circular genome 438
- 20.7 The mitochondrial genome of plants varies largely in size 442
- Further reading 449

21 Protein biosynthesis 451

- 21.1 Protein synthesis is catalysed by ribosomes 451
- 21.2 Proteins attain their three-dimensional structure by controlled folding 458
- 21.3 Nucleary encoded proteins are distributed throughout various cell compartments 464
- Further reading 470

22 Gene technology in plants 472

- 22.1 A gene is isolated 473
- 22.2 Agrobacteria have the ability to transform plant cells 483
- 22.3 Ti plasmids are used as transformation vectors 487
- 22.4 Selection of appropriate promoters enables the defined expression of an inserted gene 493
- 22.5 Genes can be turned off by transformation 494
- 22.6 Plant genetic engineering can be used for many different purposes 496
- Further reading 502

Index 505