

Contents

Chapter 1	Cell Walls and Plant Anatomy	1
Chapter 2	The Structural Polysaccharides of the Cell Wall and How They Are Studied	43
Chapter 3	Biochemistry of the Cell Wall Molecules	67
Chapter 4	Membrane Systems Involved in Cell Wall Assembly	119
Chapter 5	Biosynthesis of Cell Wall Polymers	161
Chapter 6	Principles of Cell Wall Architecture and Assembly	227
Chapter 7	The Cell Wall in Growth and Development	273
Chapter 8	Cell Walls and Plant–Microbe Interactions	319
Chapter 9	Plant Cell Walls: A Renewable Material Resource	365
	Index	411

Detailed Contents

Chapter 1 Cell Walls and Plant Anatomy	1	12. Pollen grains have a unique and sculptured protective wall.	37
A. The Derivation of Cells and Their Walls	1	13. Some plant cells have no walls.	37
1. Cells arise in specialized regions of the plant called meristems.	1	Key Terms	39
2. Walls originate in dividing cells.	3	References	39
3. Plant organ development depends on precise control of the plane of cell division and of cell expansion.	4	Chapter 2 The Structural Polysaccharides of the Cell Wall and How They Are Studied	43
4. Cytoskeletal elements predict the position of the new cross-wall before the cell divides.	8	A. Molecules of the Wall	43
5. Actin filaments help to position new cross-walls.	9	1. Primary walls isolated in a variety of ways contain the same structural polymers.	43
6. The new cross-wall must join and fuse with the mother cell wall.	10	2. Cellulose is a β -1,4-linked glucan with a disaccharide repeating unit.	44
7. A plant is constructed from two compartments: the apoplast and the symplast.	10	3. Pectins and hemicelluloses are the noncellulosic polysaccharides of primary walls.	45
B. Walls in Cell Growth and Differentiation	12	4. Wall polysaccharides are composed of 13 different sugars.	45
1. Cells become organized at an early stage into three major tissue systems.	12	5. Homogalacturonan, rhamnogalacturonan I, and rhamnogalacturonan II are three pectic polysaccharides in primary walls.	45
2. When they have stopped dividing, cells usually continue to grow in size.	16	6. Xyloglucans and arabinoxylans are the two major hemicellulosic polysaccharides in primary walls.	47
3. Depending on its position in the plant, a cell differentiates into a specific cell type.	17	7. The structural characterization of polysaccharides is difficult.	47
4. A distinction is often made between primary and secondary walls.	18	8. Cell walls contain proteins, glycoproteins, and proteoglycans that perform enzymatic, structural, defensive, and other functions.	48
5. Differentiated cell types are often characterized by functionally specialized cell walls.	20	9. Primary cell walls contain an aqueous phase.	49
C. Plant Cell Types and Their Walls	21	10. Secondary walls are composed of primary walls plus additional layers of polymers.	49
1. Shoot epidermal cells produce a waterproof cuticle that helps provide protection.	21	B. Methods for Characterizing the Structural Polysaccharides of the Cell Wall	52
2. Stomatal guard cells have asymmetrically thickened walls that allow them to reversibly change shape.	22	1. Obtaining pure polysaccharides is a prerequisite for primary structure determination.	52
3. Cells in ground tissues fill the space between epidermis and vascular tissue.	24	2. Extracellular polysaccharides secreted by cultured cells are excellent models for wall polysaccharides.	53
4. The endodermis regulates the flow of solutes through the apoplast of the root.	25	3. Enzymatic and chemical cleavage of polysaccharides has made structural studies possible.	53
5. The pericycle encircles the vascular tissue and can give rise to lateral roots.	26	4. Several methods are used for glycosyl composition analysis.	54
6. Xylem vessel elements are pipes that conduct water at negative pressure.	27	5. Glycosyl linkage compositions are determined by methylation analysis.	57
7. Phloem sieve tube elements are high-pressure tubes for conducting sugar solutions.	29	6. Glycosyl sequencing of oligo- and polysaccharides is not routine.	57
8. Transfer cells load solutes into and out of conducting elements.	31	7. Nuclear magnetic resonance (NMR) spectroscopy and mass spectrometry (MS) are important tools for determining the structures of polysaccharides.	58
9. Plasmodesmata allow adjacent cells to communicate with each other across their intervening walls.	32	8. The conformations of cell wall oligo- and polysaccharides can be investigated using NMR and computational methods.	62
10. Collenchyma and sclerenchyma provide support in young tissues.	34		
11. The shaping of mesophyll cells generates large air spaces that facilitate gaseous diffusion.	34		

9. The structure of cellulose has been determined by X-ray diffraction techniques.	63	9. AGPs, GRPs, HRGPs, and PRPs may have a common evolutionary origin.	95
Key Terms	64	F. Lignin	95
References	64	1. Lignin is a hydrophobic polymer of secondary cell walls.	95
Chapter 3 Biochemistry of the Cell Wall Molecules	67	2. Our knowledge of lignin composition is based on the chemical hydrolysis of wood.	96
A. Cellulose	67	3. Lignins in plant tissues, cells, and cell walls may be detected by histochemical staining or immunochemical localization.	99
1. β -Glucan chains hydrogen-bond together to form cellulose microfibrils.	67	4. The complexity of lignin is increased by the diversity of intermolecular linkages.	100
2. The glucan chains of cellulose microfibrils all have the same orientation.	67	5. Models for higher-order structure of lignin.	102
B. Hemicelluloses	68	6. Lignin is cross-linked to polysaccharide components in lignin-carbohydrate complexes.	103
1. Xyloglucan is a principal hemicellulose of cell walls.	68	7. Lignin composition and content vary greatly among the major groups of higher plants.	104
2. The structural features of xyloglucan have been defined by analysis of endoglucanase-released oligosaccharides.	69	8. Lignin content and composition vary in wood formed during increased mechanical stress.	105
3. Side chain substitutions on the backbone of xyloglucan affect endoglucanase cleavage of the backbone.	71	G. Suberin, Cutin, Waxes, and Silica	106
4. Arabinoxylan is the predominant hemicellulose of grasses.	72	1. Suberin is a complex hydrophobic material that forms physical and biological barriers essential for plants.	106
5. The elucidation of arabinoxylan structures has been facilitated by characterizing oligosaccharides generated by endoxylanases.	73	2. Cutin is an aliphatic polyester that is part of the plant cuticle.	110
6. Secondary walls contain xylans and glucomannans.	74	3. Epicuticular waxes form the primary interface between the outside of the plant and the aerial environment.	111
7. A galacturonic acid-containing oligosaccharide is found at the reducing end of glucuronoxylans.	74	4. Silica deposits are formed on the surface of epidermal cells as well as within the cell walls of internal tissues of many herbaceous and woody plants.	111
C. Pectic Polysaccharides	75	Key Terms	113
1. Homogalacturonans are gel-forming polysaccharides.	75	References	113
2. Rhamnogalacturonan I (RG-I) is a family of large polysaccharides with a backbone of repeating disaccharide units.	75	Chapter 4 Membrane Systems Involved in Cell Wall Assembly	119
3. Rhamnogalacturonan I (RG-I) has many arabinosyl and galactosyl residue-containing side chains.	77	A. Sites of Cell Wall Polymer Assembly in Interphase Cells	120
4. Rhamnogalacturonan II (RG-II) is the most complex polysaccharide known.	78	1. Different polymers are synthesized at different locations, in the ER and the Golgi apparatus, at the plasma membrane, or in the cell wall.	120
5. Rhamnogalacturonan II (RG-II) has an oligogalacturonide backbone.	80	2. The endoplasmic reticulum consists of a three-dimensional membrane network that extends throughout the cytoplasm and is differentiated into specialized domains.	121
6. Rhamnogalacturonan II (RG-II) has side chains with unique structures.	80	3. Golgi stacks consist of sets of flattened cisternae that are structurally distinct and exhibit a polar architecture.	122
D. Other Wall Polysaccharides	81	4. Plant Golgi stacks are dispersed throughout the cytoplasm, travel along actin filament tracks with myosin motors, and stop at ER export sites.	124
1. β -1,3/1,4-(mixed-linked) glucans are found in grasses and horsetail.	81	5. COPII vesicles bud from the ER with an external scaffold that is transferred with the vesicles to the cis side of the Golgi stacks, giving rise to the Golgi scaffold (matrix).	125
2. Enzyme-generated oligosaccharides have been used to structurally characterize mixed-linked glucans.	82	6. Intra-Golgi transport of membrane and cargo molecules is best explained by the cisternal progression/maturation model.	127
3. Callose is a β -1,3-D-glucan.	82	7. The <i>trans</i> Golgi network cisternae of plants are transient organelles that sort and package Golgi products into secretory and clathrin-coated vesicles.	128
4. Substituted galacturonans are found in some cell walls.	83	8. Maintenance of an appropriate pH in different Golgi compartments is essential for their functions.	131
5. Galactoglucomannans are found in Solanaceae cell walls and in the growth medium of Solanaceae cultured cells.	83	9. Golgi stacks are structurally and functionally differentiated in a tissue- and developmental stage-specific manner.	132
6. Some seed cell walls contain storage polysaccharides.	84	10. The process of exocytosis delivers matrix molecules to the cell wall and membrane molecules to the plasma membrane.	133
7. Gums are secreted in response to stress and are not normally considered to be wall components.	84	11. Turgor pressure has profound effects on vesicle-mediated secretion, membrane recycling from the plasma membrane, and the movement of secreted molecules through cell walls.	134
E. Proteins and Glycoproteins	85	12. Cellulose microfibrils, callose, and mixed-linked glucans are made by enzyme complexes in the plasma membrane.	136
1. Plant cell walls contain structural, enzymatic, lipid transfer, signaling, and defense proteins.	85	13. The enzymes that synthesize cellulose microfibrils in higher plants are organized into complexes that visually look like rosettes.	137
2. The repeated sequence motifs of structural proteins define their classes and their physical properties.	86		
3. Sequence motifs define proline hydroxylation, glycosylation, and intra- and intermolecular cross-linking sites.	87		
4. HRGP and PRP expression is controlled by developmental programs and triggered by external stimuli.	88		
5. Extensin-type HRGPs are rod-shaped molecules that become insolubilized by a peroxide-mediated cross-linking process.	89		
6. Arabinogalactan proteins form a family of complex proteoglycans located at the cell-surface, and in the cell wall and intercellular space.	90		
7. Classical arabinogalactan proteins are GPI-anchored proteoglycans.	92		
8. GRPs are structural cell wall proteins that participate in the assembly of vascular bundles and are present in root cap mucilage.	94		

14. Cellulose synthase complexes are pushed forward in the plasma membrane by the growing cellulose microfibrils that they extrude into the cell wall.	137	3. Processing of N-linked glycans in the plant Golgi apparatus is similar but not identical to the processing in other organisms.	178
15. Rosette complexes are concentrated in domains of the plasma membrane underlying sites of rapid cellulose synthesis.	139	4. The sites of O-linked glycosylation are defined in part by rules of proline hydroxylation.	179
16. The half-life of rosette complexes determines the length of cellulose microfibrils.	141	5. The GPI anchors of arabinogalactan proteins are both synthesized and attached to the protein backbone in the ER.	181
17. The cellulose microfibrils of primary cell walls are shorter than those of secondary cell walls.	142	C. Assembly of Polysaccharides in the Endomembrane System	182
18. Callose appears transiently during growth and development and is also formed in response to biotic and abiotic stresses.	142	1. The backbones of some complex polysaccharides are synthesized by Golgi-located enzymes encoded by genes of the cellulose synthase-like (CSL) gene family.	182
19. Callose is a transient cell wall component that is critically important for many growth and developmental processes.	143	2. Synthesis of the glucan backbone of xyloglucan in <i>trans</i> Golgi cisternae is enhanced by the cooperative assembly of glucosyl and xylosyl residues.	184
20. Stress-induced callose synthesis serves to protect cells from potentially lethal biotic and abiotic insults.	144	3. The XG-fucosyl and XG-galactosyltransferases can add fucosyl and galactosyl residues during or after synthesis of the XG backbone.	185
B. Membrane Systems Involved in <i>De Novo</i> Cell Wall Assembly during Cytokinesis	145	4. The assembly of xyloglucan occurs in <i>trans</i> Golgi and early <i>trans</i> Golgi network cisternae.	186
1. Plant Golgi stacks multiply by division.	145	5. The large number of enzymes needed to make pectic polysaccharides makes understanding their synthesis a challenge.	186
2. During mitosis and cytokinesis cytoplasmic streaming stops and the Golgi stacks redistribute to specific locations.	145	6. Synthesis of pectic polysaccharides involves enzymes localized to late <i>cis</i> , medial, and <i>trans</i> Golgi cisternae.	189
3. Somatic-type cell plate formation can be divided into four distinct phases.	147	7. Only a few steps of the pectic polysaccharide synthesis pathway have been studied biochemically <i>in vitro</i> .	189
4. Dynamin-GTPases create dumbbell-shaped, cell plate-forming vesicles containing dehydrated and possibly gelled cell wall-forming molecules.	150	8. The biosynthesis of galactomannans has parallels to the biosynthesis of xyloglucans.	191
5. The association of cisternae of the endoplasmic reticulum with forming cell plates increases over time, but the functional importance of this spatial relationship has yet to be fully explored.	152	9. Several glycosyltransferases involved in xylan biosynthesis have been identified, but the mechanism of xylan biosynthesis remains obscure.	192
6. Formation of cell walls in the syncytial endosperm and in meiocytes involves a special kind of cell plate, the syncytial-type cell plate.	153	D. Assembly of Polysaccharides at the Plasma Membrane	193
Key Terms	157	1. Studies of cellulose synthesis by <i>Gluconacetobacter xylinus</i> (<i>Acetobacter xylinum</i>) provided many of the paradigms for similar studies in higher plants.	193
References	157	2. Hydrophobic cluster analysis played a critical role in the identification of the plant <i>CESA</i> genes.	193
Chapter 5 Biosynthesis of Cell Wall Polymers	161	3. Plant <i>CESA</i> genes appear to be derived from cyanobacterial precursors.	194
A. General Mechanisms of Polymer Assembly	161	4. The cellulose synthase (<i>CESA</i>) proteins appear to correspond to the catalytic subunits of cellulose synthase complexes.	195
1. Polymers are assembled from building blocks.	161	5. The <i>CESA</i> protein is an integral protein with several domains characteristic of processive glycosyltransferases.	196
2. Nucleotide sugars are the source of the glycosyl residue building blocks used in the synthesis of cell wall polysaccharides and glycoproteins.	162	6. The Zn-binding domains of <i>CESA</i> and some CSLD proteins appear to mediate dimerization of the catalytic subunits of cellulose synthases.	197
3. Transport of nucleotide sugars into ER and Golgi cisternae is mediated by specific NDP-sugar/NMP antiporters.	166	7. The cellulose-synthesizing rosette complexes are composed of three different types of <i>CESA</i> proteins.	198
4. Wall polysaccharides are made by membrane-bound polysaccharide synthases and glycosyltransferases.	167	8. The involvement of KORRIGAN, a β -1,4-glucanase, in cellulose synthesis has yet to be proven conclusively.	198
5. Glycosyltransferases are generally nonabundant proteins.	167	9. UDP-glucose, the substrate for the cellulose and callose synthase systems, may be produced by a membrane-bound form of sucrose synthase.	199
6. All glycosidases and glycosyltransferases involved in the modification of glycoproteins and polysaccharides have a common topology and common Golgi retention mechanisms.	168	10. There are two types of callose synthase systems, a Ca^{2+} -dependent and a Ca^{2+} -independent type.	200
7. The synthesis of polysaccharide backbones involves initiation, elongation, and termination reactions.	168	11. Callose synthase is a large protein that differs in many ways from cellulose synthase.	201
8. Polysaccharide synthases can be identified biochemically by substrate and activator binding activities, by product entrapment techniques, and by catalytic activities detected in nondenaturing gels.	172	12. The ratio of tri- and tetrasaccharide units in mixed-linked glucans made <i>in vitro</i> can be altered experimentally.	202
9. The three-dimensional structure of the backbone of cell wall polysaccharides suggests synthesis by glycosyltransferases with two active sites.	172	E. Polymer Assembly in the Wall: Biosynthesis of Lignins, Waxes, Cutins, and Suberins	203
10. Polysaccharide synthesis is carefully controlled, but the control points are still poorly understood.	175	1. The secondary cell wall provides a unique environment for the polymerization of lignins from cinnamyl alcohols.	203
B. Assembly and Processing of Glycoproteins and Proteoglycans	175	2. Cinnamyl alcohols are the predominant precursors for lignin biosynthesis.	203
1. N-linked glycans of glycoproteins are assembled on dolichol lipids.	175	3. Monolignols are formed by successive enzymatic modifications of phenylalanine.	206
2. Newly synthesized N-linked glycans are transferred <i>en bloc</i> to nascent polypeptides and immediately subjected to processing by two glucosidases.	177		

4. Hydroxylation and <i>O</i> -methylation at the 3 and 5 positions on the aromatic ring are unlikely to take place at the level of cinnamic acids.	207	1. Polymers are not evenly distributed within a wall.	250
5. Coniferaldehyde is the branch point for the biosynthesis of coniferyl alcohol and sinapyl alcohol.	208	2. The middle lamella forms an adhesive boundary between adjacent cells.	252
6. Monolignols are glucosylated, stored, transported, and deglycosylated before polymerization.	208	3. Three-way junctions are rich in protein, pectin, and phenolics.	256
7. Lignin polymers are formed through enzymatic oxidation of monolignols.	210	4. The middle lamella is involved in cell separation.	257
8. Lignin polymerization is primarily due to the addition of a monolignol to a lignin polymer.	212	5. The intercellular spaces form a continuum.	258
9. Oxidative carriers or radical mediators may be involved in the polymerization process.	213	6. The expression of arabinogalactan proteins is developmentally regulated.	260
10. A nonenzymatic model for lignin polymerization has been proposed involving "dirigent" or guide proteins.	214	7. A cell wall is the product of both a cell's internal developmental program and its environmental history.	261
11. Nontraditional monomers are readily incorporated into lignin, indicating a high level of metabolic plasticity.	215	8. Wall composition and architecture, together with anatomy, contribute to the mechanical properties of plants and their products.	262
12. The deposition of lignin is both temporally and spatially controlled.	216	9. Models of the cell wall have helped us to think more clearly about their construction and function.	264
13. The composition of lignin differs between cell wall domains and cell types.	217	Key Terms	268
14. The biosynthesis of cutins, suberins, and waxes involves enzymes in chloroplasts, the endoplasmic reticulum, and cell walls.	218	References	268
Key Terms	221		
References	222	Chapter 7 The Cell Wall in Growth and Development	273
Chapter 6 Principles of Cell Wall Architecture and Assembly	227	A. Interactions Between the Cytoskeleton and the Wall	273
A. Cross-Links between Wall Polymers	227	1. Cortical microtubule reorientation changes the orientation of cellulose microfibril deposition during growth.	273
1. Wall polymers are cross-linked by covalent bonds as well as by noncovalent bonds and interactions.	227	2. Intracellular factors can alter the orientation of cortical microtubules.	276
2. Hemicelluloses bind strongly to cellulose microfibrils by noncovalent bonds.	228	3. Bundles of microtubules can define distinct domains of the wall that will thicken during cell development.	277
3. Hemicelluloses cross-link cellulose microfibrils.	229	4. The arrangement of cytoskeletal elements in one cell often relates to that in a neighboring cell.	278
4. Pectic polysaccharides are probably covalently interconnected by glycosidic bonds.	229	5. The wall is attached to receptor-like proteins in the plasma membrane.	281
5. Homogalacturonans and partially methylesterified homogalacturonans form gels.	230	B. Cell Expansion	282
6. Borate esters cross-link RG-II dimers in the wall.	231	1. A key driver of plant growth is postmitotic cell expansion.	282
7. Diferulic acids probably cross-link wall polysaccharides.	232	2. Cell expansion is usually accompanied by the deposition of new wall material.	284
8. Transesterification may produce other possible cross-links.	233	3. Wall architecture underpins anisotropic cell expansion.	287
9. Some wall proteins become insolubilized by covalent cross-links.	233	4. Dynamic remodeling of wall architecture facilitates cell expansion.	290
10. Lignin is covalently attached to hemicelluloses, and their interaction adds strength to the secondary cell wall.	234	5. Proteins that enhance wall expansion have been identified.	292
B. Architectural Principles: Putting the Polymers Together	235	6. Acidification of the wall, enhanced by auxin, may promote cell expansion.	294
1. Two coextensive polysaccharide networks underlie the structure of the primary cell wall.	235	C. Turnover, Remodeling, and Breakdown of the Wall	294
2. Walls are constructed of lamellae that are one cellulose microfibril thick.	236	1. Although plant cell walls are relatively stable compartments, many matrix polysaccharides and proteoglycans do turn over.	294
3. The primary wall is usually composed of only a small number of lamellae.	236	2. The <i>de novo</i> insertion of plasmodesmata across established walls requires wall remodeling.	296
4. The spacing of cellulose microfibrils is determined by matrix polysaccharides.	237	3. Local removal of wall material is used to create conducting elements from files of cells.	298
5. The walls of cells in which cellulose synthesis has been inhibited are composed largely of a pectin network.	240	4. Abscission of leaves and fruit is an active process that involves controlled cell separation.	299
6. The pectin network limits the porosity of the wall.	241	5. Fruit softening depends upon the expression and activity of cell wall-modifying proteins.	300
7. The cell controls the thickness of its wall.	242	6. In some seeds, wall polysaccharides can form a food reserve to be used during germination.	302
8. How proteins are integrated into wall architecture is unclear.	243	7. Lignin and suberin provide physical barriers to the turnover and degradation of secondary walls.	303
9. The orientation of newly synthesized microfibrils is determined by the cell.	244	8. Gravity sensing and mechanical stress lead to compensatory changes in cell wall synthesis and architecture.	304
10. The orientation of microfibrils within a wall may change during growth.	246	D. Cell Wall-Derived Signals in Growth and Development	306
11. The self-ordering properties of polylamellate walls can lead to high degrees of structural order.	248	1. Many signals combine to regulate plant growth.	306
12. Cell wall assembly is a hierarchical process.	249	2. Oligogalacturonides can modulate development in tobacco explants.	308
C. Architectural Variations: The Mosaic Wall	250	3. Xyloglucan-derived oligosaccharins can affect the rate of elongation growth.	308
		4. Lipo-oligosaccharides synthesized by rhizobia regulate nodule development in host plants.	310

5. Chitinases may function during normal plant development.	311
Key Terms	313
References	313

Chapter 8 Cell Walls and Plant–Microbe Interactions 319

Introduction 319

A. How Plants Detect and Respond to Microbes 320

1. Most plants are immune to most pathogens.	320
2. Signals from both pathogen and host can elicit a common defense response.	321
3. Cell surface receptors recognize common molecular patterns.	322
4. The plant's responses to danger involve a common sequence of events.	324
5. Some specialist pathogens deliver effectors to suppress the basal defense response.	326
6. Some defenses are preformed.	328
7. Defense can also be mounted at a distance.	329

B. The Wall as a Battleground 331

Pathogens enter either by force, through wounds, or through natural openings.	331
2. Plasmodesmata provide a route for pathogen spread.	333
3. Callose deposition is a local response to both pathogens and wounding.	335
4. Wall strengthening helps contain the pathogen.	336
5. Cell polarization is a common response to a pathogen.	338
6. Localized cell death restricts pathogen spread.	340
7. Cell wall fragments act as danger signals.	341
8. Plants and pathogens battle to control the release of oligogalacturonides.	343
9. PR proteins attack the walls of fungi and bacteria.	344
10. Carbohydrate-binding modules (CBMs) help enzymes attach to the wall.	345
11. Cell wall integrity is sensed by the host.	346
12. Responses to wounding and pathogens overlap.	348
13. Wall-degrading enzymes and their inhibitors coevolve.	349

C. Recycling of Cell Walls 351

1. Removing pectin is a key early step in dismantling the wall.	352
2. Lyases are important pectic enzymes for both necrotrophic pathogens and saprophytic microbes.	353
3. To fully digest hemicelluloses, enzymes act in concert.	356
4. Cellulose is tough, and its disassembly usually requires special machinery.	357
Key Terms	360
References	360

Chapter 9 Plant Cell Walls: A Renewable Material Resource 365

A. Effects of Cell Walls on the Nutritional Quality and Texture of Foods and Forage Crops 366

1. Digestibility of forage crops depends upon the properties of plant cell walls and is mainly restricted by the lignin content.	366
2. The texture of our fruits and vegetables depends to a large extent on the properties of their cell walls.	368
3. Gums, water-soluble cell wall-associated polymers, are used to stabilize emulsions and to modify the texture of processed foods and other industrial products.	369
4. Pectins are used as thickeners, texturizers, stabilizers, and emulsifiers in the food and pharmaceutical industries.	370

B. Medicinal and Physiological Properties of Cell Wall Molecules 371

1. Dietary fiber in the human diet is a residual component of plant cell walls.	371
2. The consumption of dietary fiber has been associated with benefits in human health.	372

3. Gum arabic and pectins soothe irritated or inflamed mucosal tissues.	374
4. Pectic polysaccharides from medicinal plants stimulate the immune system via Peyer's patch cells in the intestine.	374
5. Bioactive pectins from some medicinal herbs promote the proliferation of immune system cells and their activities.	375
6. Bioactive pectins inhibit tumor growth by inducing cancer cell apoptosis.	376
7. Coating of medical devices with cell wall polysaccharides alters their bioactive properties.	377

C. Cell Wall Fibers Used in Textiles 378

1. Commercially important plant fibers are derived from many types of plants and possess physical properties that are exploited in textile products.	378
2. Cotton is the most widely used plant textile fiber.	378
3. Cotton fibers are epidermal trichomes that develop on the surface of ovules.	380
4. Coconut-derived coir fibers are used extensively in tropical countries.	381
5. The physical properties of different bast fibers make them suitable for use in specialized types of textiles.	381
6. Leaf fibers are used for ropes, tea bags, currency paper, and filter-tipped cigarettes.	383

D. Wood: an Essential Product for Construction and Paper Production 384

1. Wood is one of the world's most abundant industrial raw materials.	384
2. Wood and paper properties are derived from the composition and morphology of the xylem cell walls.	385
3. The physical strength of wood depends on the multilayered structure of the wood cell walls and cell morphology.	387
4. Cell wall-degrading enzymes are increasingly used for pulp and paper processing for environmental and economic reasons.	389
5. Bark is a source of cork, tannins, waxes, and drugs.	390

E. Plant Cell Walls: the Most Important Renewable Source of Biofuels and Chemical Feedstocks 391

1. Humankind's survival may depend on the wise use of renewable natural resources.	391
2. Plant cell walls and their derivatives comprise a major fraction of the terrestrial biomass and carbon content.	394
3. Photosynthesis-driven biomass production by plants constitutes a major and sustainable energy source.	395
4. Half of the wood harvested around the world is burnt as fuel.	396
5. Producing bioethanol from lignocellulosic biomass at competitive prices will require significant investments in research.	396
6. Overcoming the lignin barrier is essential to the success of the lignocellulosic biomass-based liquid biofuel industry.	398
7. Major improvements in fermenting enzyme systems are needed to improve the efficiency of ethanol production.	400
8. New types of biorefineries are needed to produce ethanol from lignocellulosic biomass at increased efficiencies and lower cost.	400
9. Plant cell walls are a renewable source of industrial chemicals.	402
10. Large-scale biofuel and chemical feedstock production is likely to affect food prices and biodiversity.	404
11. Cell walls offer unique challenges and opportunities for genetic engineering.	405

Key Terms	407
References	407