

Contents in full

The Author	v
A Note from the Publisher on Primers in Biology	vii
Preface	ix
Acknowledgements	xii

CHAPTER 1 The Cell Cycle

1-0 Overview: Cell Reproduction	2
Cell reproduction is a fundamental feature of all living things	
Cells reproduce in discrete steps	
The ordering of cell-cycle events is governed by an independent control system	
1-1 Events of the Eukaryotic Cell Cycle	4
Chromosome duplication and segregation occur in distinct cell-cycle phases that are usually separated by gap phases	
Cytoplasmic components are duplicated throughout the cell cycle	
Cell growth is usually coordinated with cell division	
1-2 Variations in Cell-Cycle Organization	6
Cell-cycle structure varies in different cells and organisms	
Multiple rounds of chromosome duplication or segregation can occur in the same cell cycle	
The symmetry of cell division varies in different cell types	
1-3 The Cell-Cycle Control System	8
Cell-cycle events are governed by an independent control system	
The cell-cycle control system is based on oscillations in the activities of cyclin-dependent protein kinases	
Cell-cycle events are initiated at three regulatory checkpoints	
Cell-cycle progression in most cells can be blocked at checkpoints	

CHAPTER 2 Model Organisms in Cell-Cycle Analysis

2-0 Overview: Cell-Cycle Analysis in Diverse Eukaryotes	12
Mechanisms of cell-cycle control are similar in all eukaryotes	
Budding and fission yeasts provide powerful systems for the genetic analysis of eukaryotic cell-cycle control	
Early animal embryos are useful for the biochemical characterization of simple cell cycles	
Control of cell division in multicellular organisms can be dissected genetically in <i>Drosophila</i>	
Cultured cell lines provide a means of analyzing cell-cycle control in mammals	
2-1 Life Cycles of Budding and Fission Yeasts	14
Budding yeast and fission yeast divide by different mechanisms	
Yeast cells alternate between haploid and diploid states and undergo sporulation in response to starvation	
2-2 Genetic Analysis of Cell-Cycle Control in Yeast	16
Cell biological processes are readily dissected with yeast genetic methods	
Conditional mutants are used to analyze essential cell-cycle processes	
Homologous genes have different names in fission yeast and budding yeast	
2-3 The Early Embryo of <i>Xenopus laevis</i>	18
The early embryonic divisions of <i>Xenopus</i> provide a simplified system for cell-cycle analysis	
Unfertilized eggs develop from diploid oocytes by meiosis	
The early embryonic cell cycle can be reconstituted in a test tube	
2-4 The Fruit Fly <i>Drosophila melanogaster</i>	20
<i>Drosophila</i> allows genetic analysis of cell-cycle control in metazoans	
Cells of the early <i>Drosophila</i> embryo divide by a simplified cell cycle	
Gap phases are introduced in late embryogenesis	

	Adult fly structures develop from imaginal cells	
2-5	Mammalian Cell-Cycle Analysis	22
	Mammalian cell-cycle control can be analyzed in cells growing in culture	
	Mutations lead to immortalization and transformation of mammalian cells	
	Specific gene disruption is the ideal approach for assessing protein function in mammalian cells	
2-6	Methods in Cell-Cycle Analysis	24
	Cell-cycle position can be assessed by many approaches	
	Cell populations can be synchronized at specific cell-cycle stages	
	Complete understanding of cell-cycle control mechanisms requires the analysis of protein structure and enzymatic behavior	

CHAPTER 3 The Cell-Cycle Control System

3-0	Overview: The Cell-Cycle Control System	28
	The cell-cycle control system is a complex assembly of oscillating protein kinase activities	
	Multiple regulatory mechanisms govern Cdk activity during the cell cycle	
	The cell-cycle control system generates robust, switch-like and adaptable changes in Cdk activity	
3-1	Cyclin-Dependent Kinases	30
	The cyclin-dependent kinases are a small family of enzymes that require cyclin subunits for activity	
	The active site of cyclin-dependent kinases is blocked in the absence of cyclin	
3-2	Cyclins	32
	Cyclins are the key determinants of Cdk activity and can be classified in four groups	
	Cyclins contain a conserved helical core	
3-3	Control of Cdk Activity by Phosphorylation	34
	Full Cdk activity requires phosphorylation by the Cdk-activating kinase	
	Cdk function is regulated by inhibitory phosphorylation by Wee1 and dephosphorylation by Cdc25	
3-4	The Structural Basis of Cdk Activation	36
	The conformation of the Cdk active site is dramatically rearranged by cyclin binding and phosphorylation by CAK	
3-5	Substrate Targeting by Cyclin-Cdk Complexes	38
	Cyclins are specialized for particular functions	
	Cyclins can interact directly with the substrates of the associated Cdk	
	Cyclins can direct the associated Cdk to specific subcellular locations	
	Cks1 may serve as an adaptor protein that targets Cdk to phosphoproteins	
3-6	Cdk Regulation by Inhibitory Subunits	40
	Cdk inhibitors help suppress Cdk activity in G1	
	Cip/Kip proteins bind both subunits of the cyclin-Cdk complex	
	G1-Cdks are activated by Cip/Kip proteins and inhibited by INK4 proteins	
3-7	Biochemical Switches in Signaling Systems	42
	Components of the cell-cycle control system are assembled into biochemical switches	
	Switch-like behavior can be generated by various mechanisms	
	Bistability is required for an effective binary switch	
3-8	Switch-Like Activation of Cdk1	44
	Cdk1 activation at mitosis is based on positive feedback	
	Cdk switches are robust as a result of multiple partly redundant mechanisms	
3-9	Protein Degradation in Cell-Cycle Control	46
	Many cell-cycle regulators are destroyed by ubiquitin-dependent proteolysis	
	SCF catalyzes ubiquitination of phosphorylated substrates using interchangeable substrate-targeting subunits	
3-10	The Anaphase-Promoting Complex	48
	The APC initiates anaphase and mitotic exit	
	Cdc20 activates the APC in anaphase	
	APC activity is maintained in G1 by Cdh1	
	APC targets contain specific recognition sequences	

3-11	Assembling and Regulating a Cell-Cycle Oscillator	50
	Negative feedback can generate a repeating oscillator	
	Regulated braking mechanisms allow the Cdk oscillator to be paused in G1	
3-12	Transcriptional Control of Cell-Cycle Regulators	52
	A sequential program of gene expression contributes to cell-cycle control	
	Expression of a large fraction of the genes in the yeast genome is regulated during the cell cycle	
	Key gene regulatory proteins in yeast are activated at the major cell-cycle transitions	
	The E2F family controls cell-cycle-dependent changes in gene expression in metazoans	
3-13	Programming the Cell-Cycle Control System	54
	The order of cell-cycle events is determined by regulatory interactions between multiple oscillators	
	The cell-cycle control system is responsive to many external inputs	

CHAPTER 4 Chromosome Duplication

4-0	Overview: Chromosome Duplication and its Control	58
	DNA synthesis begins at replication origins	
	The cell-cycle control system activates replication origins only once in each S phase	
	Chromosome duplication requires duplication of chromatin structure	
4-1	Basic Mechanisms of DNA Synthesis	60
	The two strands of DNA are replicated by different mechanisms	
	DNA replication begins with origin unwinding and primer synthesis	
	Discontinuous DNA fragments are joined together by DNA ligase	
	Telomerase synthesizes DNA at chromosome ends	
4-2	The Replication Origin	62
	Replication origins in budding yeast contain well defined DNA sequences	
	Replication origins in animal chromosomes are defined by several factors in addition to DNA sequence	
4-3	Assembly of the Prereplicative Complex at the Replication Origin	64
	The replication origin interacts with a multisubunit protein complex	
	The ORC and accessory proteins load the Mcm helicase onto origins	
	Mcm loading involves ATP-dependent protein remodeling	
4-4	Regulation of the Prereplicative Complex	66
	Assembly of prereplicative complexes is restricted to G1 by multiple mechanisms	
	Prereplicative complex components are destroyed or inhibited in yeast as a result of Cdk activity	
	Pre-RC assembly is controlled in animals by both Cdks and the APC	
4-5	Cyclins Required for Activation of Replication Origins in Yeast	68
	Cdks and Cdc7 trigger the initiation of DNA replication	
	In budding yeast, the cyclins Clb5 and Clb6 are key activators of replication origins	
	Yeast cells lacking S cyclins can replicate their DNA	
4-6	Cyclins Required for Activation of Replication Origins in Metazoans	70
	Different cyclins control initiation of DNA replication in different stages of animal development	
	Cyclin A is a major regulator of replication initiation in cultured mammalian cells	
	DNA replication in frog embryos is triggered by cyclin E-Cdk2	
	Cyclin E-Cdk2 is a major regulator of DNA replication in <i>Drosophila</i>	
4-7	Control of Replication by the Protein Kinase Cdc7-Dbf4	72
	Cdc7 triggers the activation of replication origins	
	Cdc7 is activated during S phase by the regulatory subunit Dbf4	
	Dbf4 levels are regulated by multiple mechanisms	
4-8	Activation of the Replication Origin	74
	Replication begins with DNA unwinding at the origin	
	Late-firing origins are regulated independently	

	Replication must be completed before chromosome segregation occurs	
4-9	Basic Chromatin Structure	76
	Chromatin is complex and dynamic	
	The basic unit of chromatin structure is the nucleosome	
	Higher-order chromatin structure is also controlled by non-histone proteins, histone H1 and histone modifications	
4-10	Histone Synthesis in S phase	78
	Histone synthesis rises sharply during S phase	
	Transcription of histone genes increases in S phase	
	Histone mRNA processing and stability increase in S phase	
	The level of free histones in the cell acts as a signal to link histone synthesis to DNA synthesis	
4-11	Nucleosome Assembly on Nascent DNA	80
	Nucleosomes are distributed to both new DNA strands behind the replication fork	
	Nucleosome assembly factors load histones on nascent DNA	
4-12	Heterochromatin at Telomeres and Centromeres	82
	Heterochromatin is inherited by epigenetic mechanisms	
	Telomeres are packaged in a heritable heterochromatin structure	
	The centromere nucleates a heritable and poorly understood form of heterochromatin	
4-13	Molecular Mechanisms of Heterochromatin Duplication	84
	Duplication of heterochromatin structure involves proteins that recognize and promote localized histone modification	
	The Sir proteins form a heritable polymer at telomeres in budding yeast	
	HP1 may nucleate heritable chromatin structure at the centromere and other regions	
	Sister-chromatid cohesion in S phase prepares the cell for mitosis	

CHAPTER 5 Early Mitosis: Preparing the Chromosomes for Segregation

5-0	Overview: The Events of Mitosis	88
	The central events of mitosis are sister-chromatid separation and segregation	
	The events of early mitosis set the stage for sister-chromatid segregation	
	The completion of mitosis begins with sister-chromatid segregation	
5-1	Overview: Principles of Mitotic Regulation	90
	Phosphorylation and proteolysis control progression through mitosis	
	Mitotic events must go to completion	
	Mitotic entry and exit are major regulatory transitions with differing importance in different species	
5-2	Cyclins that Promote Mitotic Entry in Yeast	92
	Cyclin-Cdk complexes trigger mitotic entry in all eukaryotes	
	Fission yeast cells trigger mitosis with a single mitotic cyclin	
	Two pairs of mitotic cyclins control budding yeast mitosis	
5-3	Cyclins that Promote Mitotic Entry in Metazoans	94
	Mitosis in metazoans is governed by cyclins A and B	
	Vertebrate mitosis is driven by multiple forms of cyclins A and B	
	The active cyclin B1-Cdk1 complex moves from cytoplasm to nucleus in late prophase	
	Vertebrate cyclins A and B drive different mitotic events	
5-4	Regulation of Mitotic Cdks by Wee1 and Cdc25	96
	Cyclin B-Cdk1 complexes are activated rapidly in early M phase by dephosphorylation	
	Multiple Wee1-related kinases and Cdc25-related phosphatases govern Cdk1 activity in animal cells	
5-5	Switch-like Activation of Cyclin B-Cdk1 at Mitosis	98
	Mitotic Cdk1 activation involves multiple positive feedback loops	
	Cdc25B and cyclin A-Cdk help trigger cyclin B-Cdk1 activation	
5-6	Subcellular Localization of Mitotic Regulators	100
	Cyclin B1-Cdk1 is regulated by changes in its subcellular localization	

	Cyclin B1-Cdk1 location is controlled by phosphorylation of cyclin B1	
	Cdc25C localization is regulated by phosphorylation	
	Cyclin B1-Cdk1 activation and nuclear accumulation are partly interdependent	
5-7	Protein Kinases of the Polo and Aurora Families	102
	Polo-like kinases (Plks) help control spindle assembly and mitotic exit	
	Spindle function and sister-chromatid segregation are controlled in part by aurora kinases	
5-8	Preparations for Mitosis: Sister-Chromatid Cohesion	104
	Sister chromatids are held together by two mechanisms	
	Cohesin is a key mediator of sister-chromatid cohesion	
	Cohesion is established during DNA replication	
	DNA decatenation prepares sister chromatids for separation	
5-9	Entry into Mitosis: Sister-Chromatid Condensation and Resolution	106
	Chromosomes are dramatically reorganized in mitosis	
	Condensin complexes drive chromosome condensation and resolution	
5-10	Regulation of Chromosome Condensation and Resolution	108
	Mitotic Cdks act on condensin to govern the timing of chromosome condensation	
	Sister-chromatid resolution is governed by Plk and aurora B in animal cells	
 CHAPTER 6 Assembly of the Mitotic Spindle		
6-0	Overview: The Mitotic Spindle	112
	Chromosome segregation depends on the mitotic spindle	
	The mitotic spindle must be bipolar	
	Multiple mechanisms drive spindle assembly	
6-1	Microtubule Structure and Behavior	114
	Microtubules are polymers of tubulin subunits	
	Microtubules exhibit dynamic instability	
6-2	Microtubule Nucleation, Stability and Motility	116
	Cellular microtubules originate on preformed protein complexes that are usually concentrated in a microtubule-organizing center	
	Microtubule dynamics are governed by a variety of stabilizing and destabilizing proteins	
	Motor proteins move along microtubules	
6-3	The Centrosome and the Spindle Pole Body	118
	The centrosome cycle resembles the chromosome cycle	
	Centrosome behavior is determined by the centrioles	
	The yeast spindle pole body is embedded in the nuclear envelope	
6-4	Control of Centrosome Duplication	120
	Duplication of the centrosome and spindle pole body is initiated in late G1 by G1/S-Cdks	
	Centrosome duplication normally occurs once per cell cycle	
6-5	The Kinetochore	122
	The kinetochore is the major site of microtubule-chromosome attachment	
	The kinetochore provides a stable attachment to a dynamic microtubule plus end	
6-6	Early Steps in Spindle Assembly	124
	Spindle assembly begins in prophase	
	Mitotic microtubules are highly dynamic	
	Centrosome separation initiates spindle assembly	
	Centrosome maturation increases microtubule nucleation in mitosis	
6-7	Nuclear Envelope Breakdown	126
	The nuclear envelope is composed of two membranes on an underlying protein support	
	Nuclear envelope breakdown begins at nuclear pores	
	The endoplasmic reticulum and Golgi apparatus are reorganized in mitosis	
6-8	Mitotic Chromosome Function in Spindle Assembly	128
	Spindles self-organize around chromosomes	
	Microtubules can be stabilized by a gradient of Ran-GTP around chromosomes	

6-9	Attachment of Sister Chromatids to the Spindle	130
	Centrosomes search for and capture kinetochores in prometaphase	
	Some kinetochore microtubules originate at the kinetochore	
	Chromosome attachment results in tension between sister kinetochores	
6-10	Bi-Orientation of Sister Chromatids	132
	Kinetochore–microtubule attachment is stabilized by tension	
	Aurora B is required for the correction of syntelic attachments	
	Merotelic attachments are processed by multiple mechanisms	
6-11	Forces Driving Chromosome Movement	134
	Multiple forces act on chromosomes in the spindle	
	The kinetochore is a major source of poleward force	
	Microtubule flux generates poleward force	
	A polar ejection force is generated by chromosome arms	
6-12	Chromosome Congression	136
	Chromosome oscillations in prometaphase are generated by changes in the state of kinetochores	
	Microtubule flux may promote chromosome congression	

CHAPTER 7 The Completion of Mitosis

7-0	Overview: The Completion of Mitosis	140
	The final events of mitosis occur in anaphase and telophase	
	The metaphase-to-anaphase transition is initiated by ubiquitination and destruction of regulatory proteins	
	Dephosphorylation of Cdk targets drives the events of late M phase	
	APC ^{Cdc20} initiates Cdk inactivation	
7-1	Initiation of Anaphase: Activation of the APC	142
	APC ^{Cdc20} activation in early mitosis is essential for anaphase to occur	
	Phosphorylation promotes APC ^{Cdc20} activation in early mitosis	
7-2	Initiation of Anaphase: The Spindle Checkpoint	144
	Unattached kinetochores generate a signal that prevents anaphase	
	The spindle checkpoint monitors defects in microtubule attachment and kinetochore tension	
7-3	Inhibition of APC^{Cdc20} by the Spindle Checkpoint	146
	Unattached kinetochores catalyze the formation of inhibitory signaling complexes	
	The spindle checkpoint signal is rapidly turned off once kinetochores are attached	
7-4	Control of Sister-Chromatid Separation	148
	Separase is inhibited before anaphase by securin	
	In vertebrate cells Cdk1 inhibits separase by phosphorylation	
7-5	Control of Late Mitosis in Budding Yeast	150
	Cdk inactivation in mitosis in budding yeast is not due to APC ^{Cdc20} alone	
	The protein phosphatase Cdc14 is required to complete mitosis in budding yeast	
7-6	Control of Anaphase Events	152
	The anaphase spindle segregates the chromosomes	
	Dephosphorylation of Cdk targets governs anaphase spindle behavior	
7-7	Control of Telophase	154
	Dephosphorylation of Cdk substrates drives the final steps of mitosis	
	Spindle disassembly is the central event of telophase	
	Nuclear envelope assembly begins around individual chromosomes	

CHAPTER 8 Cytokinesis

8-0	Overview: Cytokinesis	158
	Cytokinesis distributes daughter nuclei into separate cells	
	Cytokinesis depends on a contractile ring and membrane deposition	
	The cleavage plane is positioned between the daughter nuclei	
	The timing of cytokinesis is coordinated with the completion of mitosis	

8-1	The Actin-Myosin Ring	160
	Bundles of actin assemble at the site of division	
	Force is generated in the contractile ring by non-muscle myosin II	
	Actin filament formation depends on formins	
8-2	Assembly and Contraction of the Actin-Myosin Ring	162
	Contractile ring function depends on accessory factors whose importance varies in different species	
	Contraction of the actin-myosin ring is regulated by activation of myosin II	
	The GTPase Rho controls actin and myosin behavior at the cleavage site	
8-3	Membrane and Cell Wall Deposition at the Division Site	164
	Membrane deposition is required during cytokinesis	
	Membrane addition occurs in parallel with actin-myosin contraction	
8-4	The Positioning and Timing of Cytokinesis in Yeast	166
	Preparations for cytokinesis in budding yeast begin in late G1	
	Fission yeast uses the nucleus to mark the division site in early mitosis	
8-5	The Positioning and Timing of Cytokinesis in Animal Cells	168
	Signals from the mitotic spindle determine the site of cleavage in animal cells	
	Multiple regulatory components at the central spindle help control cytokinesis	
	Cytokinesis is coordinated with mitosis by the spindle and Cdk1 inactivation	
8-6	Specialization of Cytokinesis in Animal Development	170
	Cytokinesis can be blocked or incomplete in some stages of development	
	Cellularization is a specialized form of cytokinesis	
8-7	Asymmetric Cell Division	172
	Asymmetric spindle positioning leads to daughter cells of unequal sizes	
	Unequal forces on the poles underlie asymmetric spindle positioning	
	The orientation of cell division is controlled by the mitotic spindle	

CHAPTER 9 Meiosis

9-0	Overview: Meiosis	176
	Sexual reproduction is based on the fusion of haploid cells	
	The meiotic program involves two rounds of chromosome segregation	
	Homologous recombination is an important feature of meiosis	
	Defects in meiosis lead to aneuploidy	
9-1	Regulation of Early Meiotic Events in Yeast	178
	The meiotic program is controlled at multiple checkpoints	
	The transcription factor Ime1 initiates the budding yeast meiotic program	
	Entry into the meiotic program is driven by the protein kinase Ime2	
9-2	Homologous Recombination in Meiosis	180
	Homologous recombination is a central feature of meiotic prophase	
9-3	Homolog Pairing in Meiotic Prophase	182
	Stages of meiotic prophase are defined by cytological landmarks	
	Homolog pairing occurs in two successive stages	
9-4	Chiasma Formation in Late Meiotic Prophase	184
	A small number of recombination sites are selected for crossover formation in zygotene	
	Crossover sites nucleate the synaptonemal complex in some species	
	Chiasmata appear in diplotene	
9-5	Controlling Entry into the First Meiotic Division	186
	Meiosis I is initiated by M-Cdk activity	
	Entry into the first meiotic division of animal cells is controlled in diplotene	
	Ndt80 and Cdk1 promote entry into the meiotic divisions of budding yeast	
	Recombination defects block entry into meiosis I	
9-6	Chromosome Attachment in Meiosis I	188
	Homolog pairs are bi-oriented on the first meiotic spindle	
	Homolog bi-orientation depends on cohesion of sister-chromatid arms	
	Homolog linkage does not involve chiasmata in some species	

9-7	Chromosome Segregation in Meiosis I	190
	Loss of sister-chromatid arm cohesion initiates anaphase I	
	The spindle checkpoint system helps control anaphase I	
	Centromeric cohesin is protected from cleavage in meiosis I	
9-8	Finishing Meiosis	192
	Meiosis I is followed by meiosis II	
	Partial Cdk1 inactivation occurs after meiosis I	
	The meiotic program is coordinated with gametogenesis	

CHAPTER 10 Control of Cell Proliferation and Growth

10-0	Overview: Control of Cell Proliferation and Growth	196
	Cell proliferation is controlled at a checkpoint in late G1	
	Progression through Start depends on an irreversible wave of Cdk activity	
	Progression through Start requires changes in gene expression	
	Cell division is often coordinated with cell growth	
10-1	Activation of Gene Expression at Start in Budding Yeast	198
	The gene regulatory proteins SBF and MBF drive expression of Start-specific genes in yeast	
	SBF and MBF are activated by Cln3-Cdk1 at Start	
	Small changes in the amount of Cln3 help trigger cell-cycle entry	
	SBF and MBF are inactivated in S phase by Clb-Cdk1 complexes	
10-2	Activation of S-Cdks in Budding Yeast	200
	G1/S-Cdks promote activation of S-Cdks	
	Multisite phosphorylation of Sic1 generates switch-like S-Cdk activation	
	G1/S- and S-Cdks collaborate to inactivate APC ^{Cdh1} after Start	
10-3	Extracellular Control of Start in Yeast: Mating Factor Signaling	202
	Yeast mating factors induce cell-cycle arrest in G1	
	Far1 has multiple functions in proliferating and arrested cells	
	Far1 phosphorylation is triggered by a G-protein signaling pathway	
10-4	Activation of Gene Expression at Start in Animals	204
	E2F transcription factors help control G1/S gene expression in animals	
	Stimulation of G1/S gene expression results from a combination of increased gene activation and decreased gene repression	
	E2F function is regulated by pRB proteins	
10-5	Regulation of E2F-pRB Complexes	206
	G1/S gene expression at Start involves the replacement of repressor E2Fs with activator E2Fs	
	Phosphorylation of pRB proteins releases E2F	
	Multiple mechanisms of E2F activation provide robust regulation of Start	
10-6	Mitogenic Signaling in Animal Cells	208
	Extracellular mitogens control the rate of cell division in animals	
	Activated mitogen receptors recruit signaling complexes to the cell membrane	
	Ras and Myc are components of many mitogenic signaling pathways	
	Activation of the PI3 kinase helps promote mitogenesis	
10-7	Activation of G1-Cdks by Mitogens	210
	Mitogenic signaling pathways lead to activation of cyclin D-Cdk complexes	
	Mitogens control cyclin D-Cdk localization and destruction	
	Mitogens and anti-mitogens control the concentrations of Cdk inhibitor proteins	
10-8	Activation of G1/S- and S-Cdk Complexes in Animal Cells	212
	G1/S-Cdk activation at Start depends on removal of the inhibitor p27	
	Cyclin A-Cdk2 activation is promoted in part by APC inhibition	
10-9	Developmental Control of Cell Proliferation	214
	Developmental signals limit cell division to specific embryonic regions	
	Embryonic divisions are limited by depletion of key cell-cycle regulators	
10-10	Overview: Coordination of Cell Division and Cell Growth	216
	Cell division and cell growth are separate processes	

	Cell growth is regulated by extracellular nutrients and growth factors	
	Cell growth and division are coordinated by multiple mechanisms	
	The size of a cell depends on its genomic content	
10-11	Control of Cell Growth	218
	Cell growth rate is determined primarily by the rate of protein synthesis	
	Extracellular nutrients and growth factors stimulate cell growth by activating the protein kinase TOR	
	TOR affects cell growth mainly by stimulating protein synthesis	
	Growth factors stimulate protein synthesis through the activation of PI3 kinase	
10-12	Coordination of Cell Growth and Division in Yeast	220
	Yeast cell growth and division are tightly coupled	
	Yeast cells monitor translation rates as an indirect indicator of cell size	
	Growth thresholds are rapidly adjustable	
10-13	Coordination of Growth and Division in Animal Cells	222
	Growth and division are coordinated by multiple mechanisms in animal cells	
	Division depends on growth in many animal cell types	
	Animal cell growth and division are sometimes controlled independently	
10-14	Control of Cell Death	224
	Animal cell numbers are determined by a balance of cell birth and death	
	Survival factors suppress the mitochondrial pathway of apoptosis	
	DNA damage and other stresses can trigger apoptosis	
 CHAPTER 11 The DNA Damage Response		
11-0	Overview: The DNA Damage Response	228
	The DNA damage response helps maintain the genome	
	ATR and ATM are conserved protein kinases at the heart of the DNA damage response	
	Replication defects trigger a DNA damage response	
11-1	Detection and Repair of DNA Damage	230
	DNA can be damaged in many ways	
	Base and nucleotide excision repair systems repair nucleotide damage	
	Double-strand breaks are repaired by two main mechanisms	
11-2	The DNA Damage Response: Recruitment of ATR and ATM	232
	ATR is required for the response to multiple forms of damage	
	ATM is specialized for the response to unprocessed double-strand breaks	
11-3	The DNA Damage Response: Adaptors and Chk1 and Chk2	234
	Protein complexes assemble at DNA damage sites to coordinate DNA repair and the damage response	
	A PCNA-like complex is required for the ATR-mediated damage response	
	Adaptor proteins link DNA damage to activation of Chk1 and Chk2	
11-4	Activation of p53 by DNA Damage	236
	p53 is responsible for long-term inhibition of cell proliferation in animal cells	
	The major regulators of p53 include Mdm2, p300 and ARF	
	Damage response kinases phosphorylate p53 and Mdm2	
11-5	Effects of DNA Damage on Progression through Start	238
	DNA damage blocks cell-cycle progression at multiple points	
	DNA damage has minor effects on progression through Start in budding yeast	
	DNA damage in vertebrate cells triggers a robust G1 arrest	
	p53 has different effects in different cell types	
11-6	Effects of DNA Damage at Replication Forks	240
	A DNA damage response is initiated at replication forks during S phase	
	ATR is the key initiator of the response to stalled replication forks	
	The DNA damage response stabilizes the replication fork	
11-7	Effects of DNA Damage on DNA Synthesis and Mitosis	242
	DNA damage in S phase blocks replication origin firing	

DNA damage blocks mitotic entry in most eukaryotes	
DNA damage blocks anaphase in budding yeast	
11-8 Responses to Mitogenic and Telomere Stress	244
Hyperproliferative signals trigger the activation of p53	
Imbalances in mitogenic stimuli promote replicative senescence in mouse cells	
Telomere degeneration promotes cell-cycle arrest in human cells	

CHAPTER 12 The Cell Cycle in Cancer

12-0 Overview: Cell-Cycle Defects in Cancer	248
Cancer cells break the communal rules of tissues	
Cancer progression is an evolutionary process driven by gene mutation	
Genetic instability accelerates cancer progression	
12-1 Gene Mutations that Drive Cancer	250
Mutations in oncogenes and tumor suppressors stimulate tumor progression	
Oncogenes can be activated by many different mechanisms	
Multiple mutations are required to cripple tumor suppressor genes	
Cancer can be initiated by mechanisms other than gene mutation	
12-2 Tissue Specificity in Cancer	252
Cancers are a complex group of diseases	
The molecular basis of tumorigenesis can vary in different tissues	
12-3 Stimulation of Cell-Cycle Entry in Cancer Cells	254
Tumor cells are independent of mitogens and resistant to anti-mitogens	
G1/S gene regulation is defective in most cancers	
Multiple mitogenic defects are required for tumor formation	
12-4 Cell Growth and Survival in Tumors	256
Cell growth is stimulated in tumors	
Tumor cells are less dependent than normal cells on survival factors	
Differentiation is often inhibited in tumor cells	
Tumor cells are resistant to the hyperproliferation stress response	
12-5 Genetic Instability in Cancer	258
Most cancer cells have unstable genomes	
Defects in the DNA damage response promote genetic instability in cancer	
Genetic instability sometimes results from an increased rate of point mutation	
Chromosomal instability is the major form of genetic instability	
12-6 Telomeres and the Structural Instability of Chromosomes	260
Defective DNA damage responses can lead to chromosomal instability	
Degenerating telomeres can lead to chromosomal instability	
12-7 Instability in Chromosome Number	262
Cancer cells often become aneuploid through a tetraploid intermediate	
Cancer cells often contain excessive numbers of centrosomes	
Mutations in mitotic spindle components contribute to chromosomal instability	
12-8 Cancer Progression	264
There are many genetic routes to a malignant cancer	
Colon cancer progression usually begins with mutations in the gene <i>APC</i>	
Two forms of genetic instability drive colorectal cancer progression	
12-9 Stopping Cancer	266
Reducing cancer mortality begins with prevention and early diagnosis	
Therapies must kill cancer cells but leave healthy cells intact	
A detailed understanding of the molecular basis of cancer may lead to rational and more specific cancer therapies	

Glossary	269
References	275
Index	287