

CONTENTS.



INTRODUCTION	Page 1
----------------------	--------

CHAPTER I

Structure of <i>Orchis</i> — Power of movement of the pollinia — Perfect adaptation of the parts in <i>Orchis pyramidalis</i> — On the insects which visit Orchids, and on the frequency of their visits — On the fertility and sterility of several Orchids — On the secretion of nectar, and on moths being purposely delayed in obtaining it	9—53
---	------

CHAPTER II.

<i>Ophreæ</i> continued — Fly and Spider <i>Ophrys</i> — Bee <i>Orphrys</i> , apparently adapted for perpetual self-fertilisation, but with paradoxical contrivances for intercrossing — The Frog Orchis; fertilisation effected by nectar secreted from two parts of the labellum — <i>Gymnadenia conopsea</i> — Greater and Lesser Butterfly Orchis; their differences and means of fertilisation — Summary on the powers of movement in the pollinia ..	54—92
--	-------

CHAPTER III.

<i>Epipactis palustris</i> ; curious shape of the labellum and its apparent importance in the fructification of the flower — <i>Cephalanthera grandiflora</i> ; rostellum aborted; early penetration of the pollentubes; case of imperfect self-fertilisation; fertilisation aided by insects — <i>Goodyera repens</i> — <i>Spiranthes autumnalis</i> ; perfect adaptation by which the pollen of a younger flower is carried to the stigma of an older flower on another plant	93—129
---	--------

CHAPTER IV

<i>Malaxis paludosa</i> ; simple means of fertilisation — <i>Listera ovata</i> ; sensitiveness of the rostellum; explosion of viscid matter; action of insects; perfect adaptation of the several organs — <i>Listera cordata</i> — <i>Neottia nidus-avis</i> ; its fertilisation effected in the same manner as in <i>Listera</i>	130—156
--	---------

CHAPTER V.

Cattleya, simple manner of fertilisation — Masdevallia, curious closed flower — Dendrobium, contrivance for self-fertilisation — Vandea, diversified structure of the pollinia; importance of the elasticity of the pedicel; its power of movement — Elasticity and strength of the caudicle — Calanthe with lateral stigmas, manner of fertilisation — Angraecum sesquipedale, wonderful length of nectary — Acropera, perplexing case, a male Orchid Page 157—210

CHAPTER VI.

Catasetidae, the most remarkable of all Orchids — The mechanism by which the pollinia of Catasetum are ejected to a distance, and are transported by insects — Sensitiveness of the horns of the rostellum — Extraordinary difference in the male, female, and hermaphrodite forms of Catasetum tridentatum — Mormodes ignea, curious structure of flower; ejection of its pollinia — Cypripedium, importance of the slipper-like form of the labellum — Secretion of nectar — Advantage derived from insects being delayed in sucking the nectar — Singular excrescences on the labellum, apparently attractive to insects 211—285

CHAPTER VII.

Homologies of Orchid-flowers — The great modification which they have undergone — Gradation of organs, of the rostellum, of the pollen-masses — Formation of the caudicle — Genealogical affinities — Mechanism of the movement of the pollinia — Uses of the petals — Production of seed — Importance of trifling details of structure — Cause of the vast diversity of structure for the same general purpose — Cause of the perfection of the contrivances in Orchids — Summary on insect-agency — Nature abhors perpetual self-fertilisation 286—360

INDEX 361—365